

1 **Title:** Faster and slower post-training recovery in futsal: multifactorial classification of
2 recovery profiles.

3
4
5 **Running head:** Classification of recovery profiles in futsal
6

7
8 **Submission type:** Original Investigation
9

10
11 **Authors:** Carolina F. Wilke^{1,2}, Felipe Augusto P. Fernandes³, Flávio Vinícius C. Martins³,
12 Anísio M. Lacerda³, Fabio Y. Nakamura^{4,5}, Samuel P. Wanner¹, Rob Duffield²
13

14 ¹ Exercise Physiology Laboratory, School of Physical Education, Physiotherapy and
15 Occupational Therapy, Universidade Federal de Minas Gerais, Belo Horizonte (MG), Brazil

16 ² Sport & Exercise Discipline Group, Faculty of Health, University of Technology Sydney,
17 Sydney (NSW), Australia

18 ³ Laboratory of Optimization and Metaheuristic Algorithms, Computer Department, Centro
19 Federal de Educação Tecnológica de Minas Gerais (CEFET), Belo Horizonte (MG), Brazil

20 ⁴ The College of Healthcare Sciences, James Cook University, Queensland, Australia

21 ⁵ Department of Medicine and Aging Sciences, “G. d’Annunzio” University of Chieti-Pescara,
22 Italy
23

24 **Corresponding Author:**

25 Carolina Franco Wilke
26

27 **Institution:** Exercise Physiology Laboratory, School of Physical Education, Physiotherapy
28 and Occupational Therapy, Universidade Federal de Minas Gerais, Belo Horizonte (MG),
29 Brazil.
30

31 **Mail address:** 6627, Av. Pres. Antônio Carlos, Campus - Pampulha - Belo Horizonte. 31270-
32 901
33

34 **Telephone:** +55 (31) 3409-2350
35

36 **E-mail:** carol_wilke@hotmail.com
37

38 **ABSTRACT**

39

40 **Purpose:** To investigate the classification of faster vs slower recovery profiles in elite futsal
41 players and factors that distinguish between them. **Methods:** Twenty-two male futsal players
42 were evaluated for the time-course of post-training recovery in countermovement jump (CMJ),
43 10m sprint, creatine kinase concentration (CK), total quality recovery (TQR) and Brunel Mood
44 Scale (fatigue and vigor) before, post, 3, 24 and 48h after a high-intensity training session.
45 Hierarchical cluster analysis was used to allocate players into different recovery profiles using
46 the area under the curve of the percentage differences from baseline of each variable. One-way
47 ANOVAs and effect sizes (ES) were used to compare the time-course of each variable and
48 players' characteristics between clusters. **Results:** Three clusters were identified and labelled
49 as faster (FR; n=6), slower physiological (SL_{phy}; n=7) and slower perceptual recovery (SL_{perc};
50 n=6), respectively. FR presented lower (better) AUC in 10m sprint than SL_{phy} (p=0.001) and
51 SL_{perc} (p=0.008). FR also showed higher (better) AUC in TQR compared to both SL_{phy}
52 (p=0.018) and SL_{perc} (p=0.026). SL_{perc} showed higher (better) AUC in CMJ than SL_{phy}
53 (p=0.014), though presented higher (worse) fatigue AUC compared to SL_{phy} (p=0.014) and FR
54 (p=0.008). AUC of CK was higher (worse) in SL_{phy} compared to FR (p=0.001) and SL_{perc}
55 (p<0.001). SL_{phy} was younger than SL_{perc} (p=0.027), whereas FR were faster 10m sprinters
56 than SL_{phy} (p=0.003) and SL_{perc} (p=0.013) and tended to have a lower VO_{2max} (ES=1.??).
57 **Conclusions:** Differing post-training recovery profiles exist in futsal players, possibly
58 influenced by their physical abilities and age/experience.

59

60 **Keywords:** Cluster analysis, classification analysis, team sport, performance.

61 **Introduction**

62 Post-exercise recovery is a complex process involving the return of performance, physiological
63 or perceptual perturbations to near pre-exercise values¹. This concept is made opaque by the
64 multi-factorial nature and varying timelines of different parameters². For example, a recent
65 meta-analysis on post-match recovery in soccer concluded that while sprint, hormonal and
66 skill/technical parameters are restored within 72 h, muscle damage, countermovement jump
67 (CMJ) and perceived well-being take longer³. However, high inter-individual variability of the
68 recovery timeline exists (i.e. faster and slower recovery); often influenced by a variety of
69 external (i.e. training/match loads, sleep and nutrition) and internal factors (i.e. aerobic and
70 intermittent-sprint capacities⁴⁻⁶), creating further challenges to interpret recovery. Thus, the
71 ability to identify faster or slower multifactorial recovery profiles may aid the prescription of
72 recovery strategies.

73

74 It is often recommended that recovery time, appropriate recovery strategies and training load
75 should be prescribed individually^{1, 7}. Albeit optimal, this invokes a challenge to coaching staff
76 given the diverse player requirements alongside restricted facilities and staff availability. In
77 this context, identifying faster and slower recovery athletes may allow practitioners to focus on
78 smaller groups based on predominant characteristics⁸. Such an approach is akin to strategies in
79 health research, where identifying certain patterns in multifaceted conditions (e.g. disease
80 diagnosis) assists professionals in selecting the most effective intervention^{9, 10}. Similar methods
81 in sport has precedent; whereby the application of a statistical classification tool to 8 screening
82 tests classified 28 professional rugby union players into 4 injury risk profiles¹¹. This
83 information was used as a basis for developing preventative programs targeting players'
84 respective needs. Accordingly, it seems reasonable to suggest that identifying athletes'
85 recovery profiles could provide means for more accurate management of recovery time,
86 interventions and training loads.

87

88 Futsal features a relatively short post-match recovery time compared to other team sports^{4, 7}.
89 Previous studies report restoration of physical performance within 24 h post-match, whereas
90 perceptual markers take longer^{12, 13}. However, these characteristics occur within the context of
91 highly congested tournaments (i.e. up to 5 games in 7 days¹⁴; and ≈ 10 training sessions per
92 microcycle¹⁵). Within such congested schedules, individuals respond and cope differently with
93 physiological and perceived fatigue, though adequate recovery is a common requirement for
94 subsequent performance. Hence, futsal constitutes an appropriate test-bed to investigate faster
95 and slower recovery profiles. Therefore, the aim of this study was to investigate whether elite
96 futsal players can be classified into different recovery profiles (i.e. faster vs slower) based on
97 multiparameter post-training evaluation. A secondary aim was to compare player
98 characteristics between recovery groups that differentiates post-training timeline
99 characteristics.

100

101

102 **Methods**

103 **Subjects**

104 Twenty-two male field futsal players participated in this study (age: 21.5 ± 5.2 years, weight:
105 69.6 ± 7.0 kg, height: 174.1 ± 5.6 cm). They were members of either the professional (PROF)
106 or under-20 (U20) squad of the same first division Brazilian team. Players provided written
107 informed consent after explanation of all procedures and were cleared by the team's medical
108 physician to participate in the study. The study was approved by the University Research Ethics
109 Committee (50166015.9.0000.5149).

110

111 **Design**

112 At the start of the 2016 pre-season, an observational design was implemented, with players
113 undertaking anthropometric and maximal aerobic capacity (VO₂max) assessments. After 1
114 (PROF) and 2 (U20) weeks of training, they underwent a high-intensity technical-tactical
115 training session representative of a typical major training session to provide a fatiguing
116 stimulus. Perceptual, physiological and performance assessments were completed before,
117 immediately and 3, 24 and 48 h post-session to evaluate the time course of recovery. In the 48h
118 preceding and prior to all experimental sessions, players were instructed to maintain their
119 habitual diet and refrain from alcohol, caffeine and high-intensity exercise.

120

121 **Methodology**

122 **Participant description**

123 At the beginning of the season, stature and body mass (in training shorts and shirt) were
124 measured. VO₂max, maximal heart rate (HRmax) and ventilatory threshold (VT) were then
125 determined during a maximal incremental test. Participants ran on a treadmill (HPX 380, Total
126 Health[®], Brazil) with fixed 1% inclination, initial speed of 6 km·h⁻¹ and continuous increments
127 of 1.0 km·h⁻¹ every minute, until volitional exhaustion. VO₂ (K4b²; Cosmed[®], Italy) and HR
128 (RS801, Polar[®], Finland) were continuously measured and recorded every minute. A 10-point
129 scale¹⁶ was used to assess their rating of perceived exertion (RPE) at the end of each stage. The
130 exercise was ceased when at least one of the following criteria was observed: the volunteer 1)
131 requested the interruption of the test; 2) failed to maintain the stipulated speed; 3) rated 10 on
132 the RPE scale; 4) showed any signs of dizziness, mental confusion, pallor, cyanosis or nausea.
133 The spirometer was calibrated before each test according to the manufacturer's instructions.
134 VO₂max and HRmax were considered the highest values measured during the test.

135

136 **Training session**

137 A 70 min high-intensity technical-tactical training session was performed in the morning on a
138 38 m x 20 m indoor futsal court. The session was developed and conducted by each squad's
139 coach to ensure ecological validity with a highly-fatiguing training session. Although sessions
140 were not explicitly standardized, coaches were instructed to be 70-min in duration of high-
141 intensity via full-court, drill-based sessions. Accordingly, both contained only futsal-specific
142 activities (i.e. small-sided games and game simulations) performed on a full court, with varying
143 technical-tactical instructions.

144

145 Before the beginning of the session, a 15-min warm up consisting of different running speeds,
146 sprints, changes of direction, and futsal specific activity was conducted by the strength and
147 conditioning coach. During the session, players were equipped with a heart rate receiver
148 (Polar[®], Finland) and a Global Positioning System device coupled with a triaxial accelerometer
149 (SPI ProX2, GPSports Systems[®], Australia). The device had a sampling frequency of 100 Hz
150 and was used in the indoor mode, whereby only the accelerometer and HR data were recorded.
151 Units were positioned between the athletes' shoulder blades in a customised designed vest.
152 Player Load was used as a measure of external training load¹⁷. Internal load was quantified
153 using HR and RPE. Mean HR was calculated relative to the players' maximal HR in the
154 incremental test (%HRmax), and the training impulse (TRIMP) according to Edwards¹⁸. RPE
155 was analysed as an indication of training intensity using the individual absolute values and as
156 an index of overall training load (session RPE; sRPE) as a product of RPE by the session
157 duration¹⁶.

158

159 **Recovery timeline characterization**

160 Players arrived 60 min prior to the session for pre-training assessments, starting with 1)
161 hydration status by urine specific gravity (USG) using a portable refractometer (Uridens Inlab,
162 São Paulo, Brazil). This was followed by 2) creatine kinase (CK) concentration from capillary
163 blood samples collected from the fingerprint (Reflotron, Roche[®], Switzerland; with intra-assay
164 coefficient of variation of <3%¹⁹). In turn, 3) players answer a customized wellness
165 questionnaire that included i) sleep hours and quality (1 = very bad and 5 = very good), ii) the
166 total quality recovery scale, ranging from 6 (worse than very, very poor recovery) to 20 (better
167 than very, very good recovery) (TQR²⁰), and iii) a Portuguese version of the Brunel Mood
168 Scale (BRUMS²¹), whereby vigor and fatigue constructs consist of the sum of four items scored
169 from 0 (nothing) to 4 (extremely).

170
171 Following the warm-up, participants performed a countermovement jump (CMJ) and 20m
172 sprint test with a 180° change of direction at 10m. The CMJ was performed on a force platform
173 (Ergo System[®], Globus, Italy) with a squat until reaching approximately 90° of knee and hip
174 flexion, followed by fast knee and hip extension, keeping the hands in the waist. The mean
175 value of four jumps separated by 15s was used for analysis. For the sprint test, photoelectric
176 cells (Multisprint, Hidrofit[®], Brazil) were positioned at the start and finish lines and at 10m
177 mark to assess time to complete 10 and 20m. Due to technological malfunction, only the first
178 10m times were used for analysis and this test is referenced as 10m sprint. Once the training
179 session was completed, CMJ, sprint and CK concentration were repeated, and approximately
180 15min later players answered to BRUMS and RPE. All procedures performed before the
181 session were then repeated 3, 24 and 48 h after training to assess the time course of recovery
182 for each variable. Due to restriction on the days of testing in the training facilities, the 24 h
183 post-training physical tests and CK concentration assessment were not performed by the PROF
184 team (n=9), though both wellness and BRUMS questionnaires were still recorded. Players were
185 familiarized with testing procedures in the days preceding the experimental session.

186 187 **Statistical analysis**

188 Firstly, to determine the profile of recovery for each marker, the percentage difference between
189 pre-training and each post-session time point was determined. These values were then
190 transformed to a z-score and used to calculate the area under the curve (AUC) of the entire 48
191 h post-training timeline for each variable via the trapezoidal method as a representation of post-
192 training recovery kinetics.

193
194 Then, using AUC of the 6 recovery parameters, an agglomerative hierarchical cluster analysis
195 based on Euclidian distance and average linkage criteria was performed (Python 2.7, Python
196 Software Foundation, <https://www.python.org/>). Briefly, each subject's data for each
197 measure is plotted in a multi-dimensional plan and the Euclidian distance between subjects is
198 calculated. The lower the distance between two subjects, the more similarities they share,
199 which further enables their classification into groups (clusters). The threshold difference of
200 115 was used to optimise clustering based initially on dendrogram differentiation, and then by
201 theoretical and meaningfulness of the resulted grouping.

202
203 Finally, to investigate the differences and characteristics between the identified clusters, the
204 AUC and the % change from baseline of each variable at each time point were compared.
205 Normality of distribution and homogeneity of variance assumptions were checked using the
206 Shapiro-Wilk and Levene's tests, respectively. Normally distributed data were compared using
207 a one-way analysis of variance, followed by the Tukey's post hoc test when applicable. Non-
208 normally distributed data were compared using Kruskal Wallis, followed by pairwise
209 comparisons when applicable (SPSS[®] software, version 22). Cohen's d effect sizes (ES) were

210 also calculated for each pairwise comparison²². The magnitudes of the ES were qualitatively
211 interpreted using the following thresholds: < 0.2, trivial; 0.2–0.6, small; 0.6–1.2, moderate;
212 1.2–2.0, large; 2.0–4.0, very large and; > 4.0, nearly perfect ²².
213 .
214

215 **Results**

216 Cluster analysis resulted in the classification of players into 3 respective groups (Figure 1A).
217 Cluster 1 consisted of 6 U20 players, cluster 2 included 7 players (4 U20 and 3 PROF), and 6
218 players were grouped in cluster 3 (2 U20 and 4 PROF). However, 3 athletes were not included
219 in any group due to the average linkage distance threshold (1 U20 and 2 PROF).

220

221 As context to the recovery profiles, both external and internal training loads were not
222 significantly different ($p>0.05$) between the 3 groups (Table 1); however, small - moderate
223 effect sizes were evident for PL and TRIMP between clusters 2 and 3 ($ES = -0.95$, $CI = [-2.10-$
224 $0.21]$ and $-1.07 [-2.22- 0.09]$, respectively); as well as for % HRmax between clusters 1 and 2
225 ($-0.86 [-2.10- 0.37]$).

226

227

* Table 1 about here *

228

229 Figure 1B presents the AUC for each recovery variable of the respective clusters. Of note,
230 lower AUC for 10m sprint, CK and Fatigue; and higher AUC for CMJ, TQR and Vigor
231 represents a better post-session response and/or a shorter time to return to baseline. For ease of
232 interpretation, clusters with the best or worse AUC in each variable will be reported to contrast
233 with other clusters. Cluster 3 showed significantly higher (better) AUC in CMJ than cluster 1
234 ($p=0.014$; $ES=1.63$, $CI = [0.65- 2.60]$). For 10m sprint performance, AUC of cluster 2 was
235 significantly lower (better) than clusters 1 ($p=0.001$; $-1.82 [-2.79- -0.86]$) and 3 ($p=0.008$; $-$
236 $2.59 [-3.54- -1.64]$). A significantly higher (worse) AUC of CK was evident in cluster 1
237 compared to clusters 2 and 3 ($p=0.001$; $2.26 [1.33- 3.20]$ and $p<0.001$; $3.46 [2.49- 4.43]$,
238 respectively). Cluster 2 showed higher (better) AUC in TQR compared to both cluster 1
239 ($p=0.018$; $1.43 [0.49- 2.36]$) and cluster 3 ($p=0.026$; $1.55 [0.63- 2.46]$). Similarly, AUC for
240 vigor scores in cluster 2 was significantly higher (better) than cluster 3 ($p=0.003$; $2.07 [1.15-$
241 $2.99]$). Regarding fatigue, cluster 3 presented significantly higher (worse) AUC compared to
242 cluster 1 ($p=0.014$; $1.50 [0.53- 2.47]$) and cluster 2 ($p=0.008$; $1.69 [0.72- 2.66]$). Collectively,
243 based on the most prominent characteristics of recovery depicted by each cluster, we classified
244 them as follows:

245 Cluster 1 = slower physiological recovery group (SL_{phy})

246 Cluster 2 = faster recovery group (FR)

247 Cluster 3 = slower perceptual recovery group (SL_{perc}).

248

249

* Figure 1 about here *

250

251 Subsequently, to test the appropriateness of the above cluster descriptors, the mean percentage
252 changes relative to baseline in each parameter over the 48 h post-training recovery were
253 compared (Figure 2). Immediately post-session, changes in physical performance (CMJ and
254 10m sprint) and CK were not significantly different between the 3 clusters (CMJ: $p=0.467$;
255 10m sprint: $p=0.692$; CK: $p=0.447$; ES ranging from $-0.60 [-1.60- 0.41]$ to $0.71 [-0.20- 1.62]$).
256 However, CK concentration presented a significantly higher increase in SL_{phy} at 3 h post-
257 session compared to FR ($p=0.027$; $1.27 [0.28- 2.26]$) and SL_{perc} ($p=0.022$; $1.35 [0.37- 2.34]$),
258 as well as higher changes 48 h after training than SL_{perc} ($p=0.005$; $2.61 [1.62- 3.60]$). The %
259 change in 10m sprint performance of FR participants was significantly lower (better) compared
260 to SL_{phy} at 3 h ($p<0.001$; $-2.81 [-3.74- -1.88]$) and 48 h ($p=0.007$; $-1.85 [-2.79- -0.91]$); as well
261 as lower than SL_{perc} players 3 h ($p=0.002$; $-2.07 [3.01- -1.13]$). Contrastingly, 3 h after training
262 the changes in CMJ were significantly better in the SL_{perc} group compared to FR ($p=0.013$;
263 $1.59 [0.63- 2.55]$) and SL_{phy} ($p=0.001$; $2.16 [1.19- 3.13]$), whereas differences were not
264 significant at 48 h.

265
266 In respect to perceptual responses, no significant differences amongst clusters were evident in
267 the change in TQR 3h post-session ($p=0.246$). However, its subsequent increase was
268 significantly higher in FR compared to SL_{per} at 24h ($p=0.041$; 1.12 [0.19- 2.05]) and compared
269 to SL_{phy} at 48h post-session ($p=0.027$; 1.37 [0.40- 2.34]). Similarly, the decrease in vigor
270 immediately ($p=0.218$) and 3h after the session ($p=0.245$) were not significantly different
271 between clusters. However, changes were significantly higher in FR compared to SL_{perc} at both
272 24h ($p=0.011$; 1.88 [0.96 to 2.80]) and 48h after training ($p=0.012$; 2.07 [1.16-2.98]). Fatigue
273 scores were only different 24h post-session, when the SL_{perc} group presented higher changes
274 from baseline compared to SL_{phy} ($p=0.011$; 1.88 [0.89 to 2.87]).

275
276 * Figure 2 about here *

277
278 When comparing participant characteristics between the 3 clusters (Table 3), anthropometric
279 measures were not significantly different ($p>0.05$), though SL_{perc} players were younger than
280 SL_{phy} ($p=0.027$; -1.04 [-2.03- -0.05]) and moderate effect sizes were evident compared to FR
281 ($p=1.000$; -0.55 [-1.52- 0.42]) clusters. Regarding physical performance, SL_{phy} and SL_{perc}
282 players were significantly faster in the 10m sprint compared to FR ($p=0.003$; -1.99 [-2.96- -
283 1.02] and $p=0.013$; -1.89 [-2.84- 0.93], respectively). Although no significant difference was
284 evident for VO_{2max} ($p=0.128$), there was a moderate - large effect for higher values in FR in
285 comparison to SL_{phy} (1.13 [0.15- 2.11]) and for SL_{perc} in comparison to SL_{phy} (0.70 [-0.33-
286 1.73]).

287
288 From baseline measures, only vigor scores were significantly higher in SL_{phy} than in FR
289 participants ($p=0.041$, 1.16 [0.23- 2.10]). Moderate effect sizes were found for TQR (-1.16 and
290 -0.88), vigor (-1.59 and -1.05) and sleep quality (-0.83 and 1.14) when comparing FR to both
291 SL_{phy} and SL_{perc} , respectively. In addition, effect sizes were moderate for tension (-0.81 and -
292 0.84) and depression (-0.98 and -0.63) when comparing SL_{perc} to FR and SL_{phy} , respectively.

293
294 * Table 2 about here *

295 296 297 **Discussion**

298 This study investigated the identification of faster and slower post-training recovery profiles in
299 elite futsal players, and the distinguishing characteristics between respective groups. The
300 cluster analysis differentiated 3 groups based on 6 recovery parameters (cluster 1 = SL_{phy} ;
301 cluster 2 = FR; cluster 3 = SL_{perc}). FR players demonstrated better post-training recovery in 4
302 of the 6 measures (10m sprint, TQR, vigor, fatigue), showed slower sprint performance and
303 moderate effects for increased VO_{2max} . SL_{phy} players showed poorer sprint performance and
304 higher CK concentrations, despite a tendency to report better perceived recovery (TQR, vigor
305 and fatigue). Conversely, SL_{perc} players were older than SL_{phy} , and reported poorer mood states
306 (vigor and fatigue) despite no overt decrement in any physical performance. Consequently, a
307 multi-parameter classification of recovery state may be possible to differentiate recovery
308 characteristics and guide training and recovery practices.

309
310 Given the technical-tactical nature of the session replicating ecologically valid high-intensity
311 training routines, training load was not precisely standardized for all players. However, despite

312 better pre-training TQR of FR players, no differences in training load parameters PL, %HRmax
313 and RPE were between clusters (Table 1). Aligned with these results, comparisons of post-
314 session CMJ, 10m sprint and CK changes from baseline were not significantly different
315 between clusters, supporting previously reported association between training loads and
316 physical performance after a soccer match²³. Therefore, it is reasonable to infer that factors
317 aside from training loads would explain the distinct recovery profiles.

318

319 Discussing the respective groups in isolation, FR demonstrated faster recovery in 10m sprint,
320 TQR, vigor and fatigue than the other groups. We propose this represents a “preferred”
321 recovery profile given reduced extent of post-training fatigue or faster return to pre-testing is
322 considered optimal^{1, 4}. Additionally, the aligned response of objective and subjective
323 parameters agrees with integrative models of fatigue²⁴, supporting recent perspectives of the
324 mechanisms underpinning recovery². Interestingly, defining characteristics of this FR cluster
325 were the slowest 10m time compared to the other clusters and a tendency (moderate ES)
326 towards higher VO_{2max} compared to the other two clusters. Such a finding aligns with previous
327 research reporting that players with higher YoYo performance showed faster post-match
328 recovery following a rugby league match than their counterparts with lower performance⁴.
329 Accordingly, the profile of futsal players who may be considered to have better “recovery
330 capability” may relate to higher aerobic fitness. However, the tendency towards lower %HR_{max}
331 during the session for FR players compared to SL_{phy} groups raises the question of whether
332 physical capacity or training load may best explain the difference in recovery profile .

333

334 SL_{phy} players exhibited the worst AUC for CMJ and CK, based on a decrease in CMJ 3h post-
335 session and the sustained increase in CK up to 48h. This profile represents higher peaks in
336 muscle damage and reduced power during the 48h post-training, which could risk optimal
337 performance at ensuing training/competition sessions during congested schedules, and
338 represent the most important group to intervene to aid recovery¹. Notably, SL_{phy} presented
339 faster 10m sprint time before training, as well as a tendency (moderate ES) towards lower
340 VO_{2max} than FR. In this case, it is not unexpected that high power athletes with higher
341 proportion of fast-twitch muscle fibers may experience greater decrease in power performance
342 and longer time for muscle damage repair^{25, 26}. Albeit speculative, this rationale also aligns
343 with the greater decreases in speed previously observed in faster futsal players after a
344 preseason²⁷. Accordingly, extra attention to the neuromuscular recovery status of high
345 speed/power athletes during congested schedules can be beneficial.

346

347 The SL_{perc} group reported worse fatigue and vigor AUC, representative of worse scores relative
348 to baseline 24h and 48 h after the session. However, these players also depicted better CMJ
349 and CK recovery profiles. These results contradicts our expectations of an overall slower
350 recovery profile, and is likely to represent differences often reported by practitioners between
351 an athlete’s perception of recovery and the observed physical performance in a session²⁸. The
352 environmental or psycho-physiological factors that affect these perceptions remain speculative,
353 but this profile highlights perception of recovery as an important factor to consider in sub-
354 groups of players. Given these players were older than SL_{phy} participants, it is possible that age
355 and experience affected players’ perceptual mood/recovery contributing to the observed
356 mismatch between objective and subjective parameters’ timeline of recovery in SL_{perc} and SL_{phy}

357 clusters. As evidence, years of experience in professional Australian football have been
358 associated with higher RPE for a constant external training load²⁹.

359

360 Despite the attempt to classify and explain recovery clusters, several limitations need to be
361 further acknowledged. To partially overcome the restricted number of players constituting a
362 futsal team, we evaluated two age/skill level groups in separate sessions; albeit it appears that
363 training load *per se* was not the determinant of the different recovery profiles, the influence on
364 the current findings remains uncertain. We also acknowledge that sample size can still restrict
365 the extrapolation of our findings, as well as the limitation of the physiological dimension to a
366 single muscle damage marker (CK). Moreover, it is important to address that 3 players were
367 not nested to any cluster, showing that not all athletes fit in a general classification of recovery,
368 and therefore the use of this technique to guide training loads and recovery practices can be
369 limited. Finally, we recognise that this study represents responses to one session and the
370 methodological assessment may not be practical to high performance teams.

371

372 **Practical Applications**

373 Given the distinct timeline of recovery of physical, physiological, perceptual and mood
374 markers, recovery monitoring should include both objective and subjective measures,
375 alongside training load measures to aid appropriate interpretation. Based on such multifactorial
376 recovery timeline, our results provide initial insights to the use of statistical tools as a diagnostic
377 approach, discriminating smaller groups within a team to support the prescription of training
378 and recovery according to main individual needs. Future studies are thus encouraged to adapt
379 more functional approaches for recovery profile assessment.

380

381 **Conclusions**

382 Differing post-training recovery profiles were evident in futsal players. A faster global
383 (physical and psychological) recovery profile existed, possibly positively affected by higher
384 aerobic capacity. Interestingly, two groups were classified with distinct slower recovery
385 profiles conditioned by responses in either physiological or perceptual parameters, potentially
386 influenced by higher speed/power performance and higher age/experience of athletes,
387 respectively.

388

389

390 **ACKNOWLEDGMENTS**

391 This investigation was supported by Fundação de Amparo à Pesquisa de Minas Gerais
392 (FAPEMIG; APQ-02953-16), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior
393 (CAPES), and the Brazilian Ministry of Sport. The authors are grateful to Minas Tennis Clube
394 for allowing the development of this research, the coaching staff for having embraced the
395 project and the players for their participation.
396

397 **References**

398

- 399 1. Kellmann M, Bertollo M, Bosquet L, et al. Recovery and Performance in Sport:
400 Consensus Statement. *Int J Sports Physiol Perform.* 2018;13(2):240-245.
- 401 2. Minett GM, Duffield R. Is recovery driven by central or peripheral factors? A role for
402 the brain in recovery following intermittent-sprint exercise. *Front Physiol.* 2014;5:24.
- 403 3. Silva JR, Rumpf MC, Hertzog M, et al. Acute and Residual Soccer Match-Related
404 Fatigue: A Systematic Review and Meta-analysis. *Sports Med.* 2018;48(3):539-583.
- 405 4. Johnston RD, Gabbett TJ, Jenkins DG, Hulin BT. Influence of physical qualities on
406 post-match fatigue in rugby league players. *J Sci Med Sport.* 2015;18(2):209-213.
- 407 5. Skein M, Duffield R, Minett GM, Snape A, Murphy A. The effect of overnight sleep
408 deprivation after competitive rugby league matches on postmatch physiological and
409 perceptual recovery. *Int J Sports Physiol Perform.* 2013;8(5):556-564.
- 410 6. Wanner SP, Wilke CF, Duffield R. Nutritional strategies for maximizing recovery from
411 strenuous exercise in the heat: An important role for carbohydrate (sago)
412 supplementation. *Temperature (Austin, Tex.).* 2016;3(3):366-368.
- 413 7. Nedelec M, McCall A, Carling C, Legall F, Berthoin S, Dupont G. Recovery in soccer:
414 part I - post-match fatigue and time course of recovery. *Sports Med.* 2012;42(12):997-
415 1015.
- 416 8. Doeven SH, Brink MS, Kosse SJ, Lemmink K. Postmatch recovery of physical
417 performance and biochemical markers in team ball sports: a systematic review. *BMJ
418 open sport & exercise medicine.* 2018;4(1):e000264.
- 419 9. Guo Q, Lu X, Gao Y, et al. Cluster analysis: a new approach for identification of
420 underlying risk factors for coronary artery disease in essential hypertensive patients.
421 *Sci Rep.* 2017;7:43965.
- 422 10. Shannon W. Cluster analysis *Epidemiology and Medical Statistics.* Vol 27. 1 ed:
423 Elsevier; 2007:342-366.
- 424 11. Evans KL, Hughes J, Williams MD. Reduced severity of lumbo-pelvic-hip injuries in
425 professional Rugby Union players following tailored preventative programmes. *J Sci
426 Med Sport.* 2018;21(3):274-279.
- 427 12. Moreira A, Costa EC, Coutts AJ, Nakamura FY, Silva DAd, Aoki MS. Cold water
428 immersion did not accelerate recovery after a futsal match. *Revista Brasileira de
429 Medicina do Esporte.* 2015;21:40-43.
- 430 13. Tessitore A, Meeusen R, Pagano R, Benvenuti C, Tiberi M, Capranica L. Effectiveness
431 of active versus passive recovery strategies after futsal games. *J Strength Cond Res.*
432 2008;22(5):1402-1412.
- 433 14. Charlot K, Zongo P, Leicht AS, Hue O, Galy O. Intensity, recovery kinetics and well-
434 being indices are not altered during an official FIFA futsal tournament in Oceanian
435 players. *J Sports Sci.* 2016;34(4):379-388.
- 436 15. Soares-Caldeira LF, de Souza EA, de Freitas VH, de Moraes SM, Leicht AS, Nakamura
437 FY. Effects of additional repeated sprint training during preseason on performance,
438 heart rate variability, and stress symptoms in futsal players: a randomized controlled
439 trial. *J Strength Cond Res.* 2014;28(10):2815-2826.
- 440 16. Foster C, Daines E, Hector L, Snyder AC, Welsh R. Athletic performance in relation to
441 training load. *Wis Med J.* 1996;95(6):370-374.
- 442 17. Boyd LJ, Ball K, Aughey RJ. The reliability of MinimaxX accelerometers for
443 measuring physical activity in Australian football. *Int J Sports Physiol Perform.*
444 2011;6(3):311-321.
- 445 18. Edwards S. *The Heart Rate Monitor Book:* Polar CIC; 1993.

- 446 **19.** Howatson G, Milak A. Exercise-induced muscle damage following a bout of sport
447 specific repeated sprints. *J Strength Cond Res.* 2009;23(8):2419-2424.
- 448 **20.** Kentta G, Hassmen P. Overtraining and recovery. A conceptual model. *Sports Med.*
449 1998;26(1):1-16.
- 450 **21.** Rohlf's ICPdM, Rotta TM, Luft CDB, Andrade A, Krebs RJ, Carvalho Td. A Escala de
451 Humor de Brunel (Brums): instrumento para detecção precoce da síndrome do excesso
452 de treinamento. *Revista Brasileira de Medicina do Esporte.* 2008;14:176-181.
- 453 **22.** Hopkins WG. A Scale of Magnitudes for Effect Statistics 2002. 2002. Accessed May
454 5, 2018.
- 455 **23.** Rowell AE, Aughey RJ, Hopkins WG, Stewart AM, Cormack SJ. Identification of
456 Sensitive Measures of Recovery After External Load From Football Match Play. *Int J*
457 *Sports Physiol Perform.* 2017;12(7):969-976.
- 458 **24.** St Clair Gibson A, Swart J, Tucker R. The interaction of psychological and
459 physiological homeostatic drives and role of general control principles in the regulation
460 of physiological systems, exercise and the fatigue process - The Integrative Governor
461 theory. *Eur J Sport Sci.* 2018;18(1):25-36.
- 462 **25.** Pimenta EM, Coelho DB, Cruz IR, et al. The ACTN3 genotype in soccer players in
463 response to acute eccentric training. *Eur J Appl Physiol.* 2012;112(4):1495-1503.
- 464 **26.** Pimenta EM, Coelho DB, Veneroso CE, et al. Effect of ACTN3 gene on strength and
465 endurance in soccer players. *J Strength Cond Res.* 2013;27(12):3286-3292.
- 466 **27.** Nakamura FY, Pereira LA, Rabelo FN, Ramirez-Campillo R, Loturco I. Faster Futsal
467 Players Perceive Higher Training Loads and Present Greater Decreases in Sprinting
468 Speed During the Preseason. *J Strength Cond Res.* 2016;30(6):1553-1562.
- 469 **28.** Doeven SH, Brink MS, Frencken WGP, Lemmink K. Impaired Player-Coach
470 Perceptions of Exertion and Recovery During Match Congestion. *Int J Sports Physiol*
471 *Perform.* 2017;12(9):1151-1156.
- 472 **29.** Gallo T, Cormack S, Gabbett T, Williams M, Lorenzen C. Characteristics impacting on
473 session rating of perceived exertion training load in Australian footballers. *J Sports Sci.*
474 2015;33(5):467-475.

475

476

477 **FIGURE CAPTIONS**

478

479 **Figure 1.** **A)** Dendrogram resulted from the cluster analysis. **B)** Area under the curve (AUC)
480 of each recovery variable for the 3 clusters. The data in panel B is expressed as mean \pm SD. ;
481 A = different from Cluster 1; B = different from Cluster 2. Legend: CMJ: countermovement
482 jump, CK: creatine kinase, TQR: total quality recovery scale.

483

484 **Figure 2.** Percentage difference from baseline obtained at each time point (post-training, 3 h,
485 24 h and 48 h hours post training) of the 3 clusters in each recovery parameter. a)
486 countermovement jump (CMJ), b) 10m sprint, c) creatine kinase (CK), d) total quality recovery
487 (TQR) scale, e) Vigor, e) FatigueA = different from SL_{phy} ; B = different from FR; C = different
488 from SL_{perc} .

489 **Tables**

490

491 **Table 1:** Training load parameters of the three clusters (mean \pm SD).

492

Training load parameter	Cluster 1	Cluster 2	Cluster 3	p	ES
Player Load	596 \pm 94	536 \pm 113	652 \pm 104	0.292	-0.50 / -0.95 / 0.51
% HRmax	81 \pm 5%	77 \pm 4%	79 \pm 4%	0.343	-0.86 / -0.54 / -0.44
TRIMP	228 \pm 29	215 \pm 22	242 \pm 22	0.301	-0.42 / -1.07 / 0.49
RPE	5.8 \pm 1.3	6.3 \pm 2.0	7.0 \pm 1.7	0.502	0.25 / -0.36 / 0.71
sRPE	397 \pm 83	446 \pm 137	503 \pm 110	0.353	0.25 / -0.36 / 0.71

493

494

495 ES = Effect size, presented in the following order of comparisons: Cluster 1 vs Cluster 2 /
496 Cluster 2 vs Cluster 3 / Cluster 1 vs Cluster 3. B = different from Cluster 2.

497

498 **Table 2:** Age, anthropometry, physical performance and pre-training measures of the three
 499 clusters (mean \pm SD).
 500

	Cluster 1 (SLphy)	Cluster 2 (FR)	Cluster 3 (SLperc)	p	ES
Age / Anthropometry					
Age (years)	18.3 \pm 1.0	20.8 \pm 3.4	24.0 \pm 6.5 ^A	0.027	0.89 / -0.55 / 1.03
Body mass (kg)	68.2 \pm 10.8	70.0 \pm 3.2	70.4 \pm 6.1	0.857	0.19 / -0.07 / 0.22
Stature (cm)	174.2 \pm 7.1	175.1 \pm 7.0	172.7 \pm 3.4	0.778	0.12 / 0.40 / -0.24
Physical performance					
VO ₂ max (mlO ₂ .kg ⁻¹ .min ⁻¹)	48.9 \pm 4.0	54.2 \pm 4.5	51.9 \pm 3.6	0.128	1.13 / 0.52 / 0.70
%VO ₂ max at VT	43.3 \pm 4%	45.5 \pm 12%	52.2 \pm 14%	0.466	0.23 / -0.49 / 0.76
CMJ (cm)	33.7 \pm 4.2	32.7 \pm 4.3	30.9 \pm 1.4	0.407	-0.22 / 0.51 / -0.79
Sprint 0-10m (s)	1.53 \pm 0.06	1.64 \pm 0.03 ^A	1.55 \pm 0.05 ^B	0.002	1.99 / 1.89 / 0.33
Pre-training measures					
CK (U/L)	198 \pm 129	168 \pm 89	327 \pm 370	0.908	-0.25 / -0.51 / 0.41
TQR	14.7 \pm 1.4	13.0 \pm 1.3	14.3 \pm 1.5	0.099	-1.16 / -0.88 / -0.21
Vigor	11.5 \pm 1.8	7.7 \pm 2.6 ^A	10.5 \pm 2.3	0.035	-1.59 / -1.05 / -0.44
Fatigue	2.8 \pm 2.7	4.6 \pm 2.4	1.8 \pm 1.2	0.105	0.63 / 1.33 / -0.42
Tension	3.0 \pm 2.5	2.9 \pm 2.5	1.2 \pm 1.0	0.279	-0.05 / 0.81 / -0.84
Depression	0.5 \pm 0.5	1.0 \pm 1.0	0.2 \pm 0.4	0.142	0.57 / 0.98 / -0.63
Anger	0.7 \pm 1.6	1.1 \pm 2.2	0.7 \pm 1.6	0.867	0.23 / 0.23 / 0.00
Confusion	1.0 \pm 1.5	1.7 \pm 2.2	0.7 \pm 1.2	0.552	0.35 / 0.54 / -0.22
Urine specific gravity	1020 \pm 7	1026 \pm 7	1026 \pm 7	0.321	0.78 / 0.07 / 0.71
Sleep hours	7.5 \pm 0.9	7.0 \pm 0.9	6.7 \pm 1.1	0.387	-0.46 / 0.34 / -0.73
Sleep quality	3.5 \pm 0.5	3.0 \pm 0.6	3.8 \pm 0.8	0.100	-0.83 / -1.14 / 0.46

501
 502 ES = Effect size, presented in the following order of comparisons: Cluster 1 vs Cluster 2 /
 503 Cluster 2 vs Cluster 3 / Cluster 1 vs Cluster 3. SL_{phy} = slower physiological recovery, FR =
 504 faster recovery, SL_{perc} = slower perceptual recovery, CK = creatine kinase, CMJ =
 505 countermovement jump, TQR = total quality recovery scale, VO₂max = maximal oxygen
 506 consumption, %VO₂max at VT = percentage of maximal oxygen consumption at the time the
 507 ventilatory threshold was reached. A = different from Cluster 1 (SL_{phy}); B = different from
 508 Cluster 2 (FR).

509
 510
 511
 512
 513