

**Factors governing microalgae harvesting efficiency by flocculation using cationic  
polymers**

**Submitted to: *Bioresource Technology***

**July 2021**

Hang P. Vu<sup>a</sup>, Luong N. Nguyen<sup>a</sup>, Benjamin Emmerton<sup>b</sup>, Qilin Wang<sup>a</sup>, Peter J. Ralph<sup>b</sup> and  
Long D. Nghiem<sup>a,c,\*</sup>

<sup>a</sup> Center for Technology in Water and Wastewater, School of Civil and Environmental  
Engineering, University of Technology Sydney, Sydney, NSW 2007, Australia

<sup>b</sup> Climate Change Cluster, Faculty of Science, University of Technology Sydney, Sydney,  
NSW 2007, Australia


<sup>c</sup> Nguyen Tat Thanh University, NTT Institute of Hi-Technology, Ho Chi Minh City,  
Vietnam

\*Corresponding author:


Long D. Nghiem: Centre for Technology in Water and Wastewater, School of Civil and  
Environmental Engineering, University of Technology Sydney, NSW 2007, Australia

Phone: +61 2 95142625 E-mail: [duclong.nghiem@uts.edu.au](mailto:duclong.nghiem@uts.edu.au)

## Supplementary data



**Figure 1S:** Three step cultivation of *Chlorella vulgaris*



**Figure 2S:** Microscopic images of *Chlorella vulgaris* cells (A) before (A) and (B) after (B) flocculation with cationic polymer.