

# An Examination of the Association Between Subjective Distress and Functional Connectivity During Discarding Decisions in Hoarding Disorder

Hannah C. Levy, Andrew Poppe, Jaryd Hiser, Bethany M. Wootton, Lauren S. Hallion, David F. Tolin, and Michael C. Stevens

## ABSTRACT

**BACKGROUND:** Individuals with hoarding disorder (HD) demonstrate exaggerated subjective distress and hyperactivation of cingulate and insular cortex regions when discarding personal possessions. No prior study has sought to determine whether this subjective distress is associated with specific profiles of abnormal brain function in individuals with HD.

**METHODS:** We used multimodal canonical correlation analysis plus joint independent component analysis to test whether five hoarding-relevant domains of subjective distress when deciding to discard possessions (anxiety, sadness, monetary value, importance, and sentimental attachment) are associated with functional magnetic resonance imaging-measured whole-brain functional connectivity in 72 participants with HD and 44 healthy controls.

**RESULTS:** Three extracted components differed between HD participants and healthy control subjects. Each of these components depicted an abnormal profile of functional connectivity in HD participants relative to control subjects during discarding decisions, and a specific distress response profile. One component pair showed a relationship between anxiety ratings during discarding decisions and connectivity among the pallidum, perirhinal ectorhinal cortex, and dorsolateral prefrontal cortex. Another component comprised sadness ratings during discarding decisions and connectivity in the pallidum, nucleus accumbens, amygdala, and dorsolateral prefrontal cortex. The third component linked HD brain connectivity in several dorsolateral prefrontal cortex regions with perceived importance ratings during discarding decisions.

**CONCLUSIONS:** The findings indicate that in patients with HD, the subjective intensity of anxiety, sadness, and perceived possession importance is related to abnormal functional connectivity in key frontal and emotional processing brain regions. The findings are discussed in terms of emerging neurobiological models of HD.

<https://doi.org/10.1016/j.bpsc.2020.12.014>

Hoarding disorder (HD) is characterized by difficulty discarding possessions, resulting in excessive clutter in the home that precludes the normal use of living spaces (1). Neuroimaging studies indicate that patients with HD have characteristic functional abnormalities in cingulate and insular cortex regions elicited during symptom provocation tasks (2,3). Specifically, when patients with HD make decisions to discard possessions (either real or imagined personal possessions), they exhibit relatively greater activation of the anterior cingulate cortex (ACC) and insula. In contrast, personally irrelevant, emotionally neutral decisions result in activity levels in these brain regions that are generally blunted compared to nonpatient controls or patients with obsessive-compulsive disorder (2,3). Furthermore, neural activity in the ACC and insula correlated with self-reported hoarding severity in these studies (2,3). Although other HD brain activation abnormalities have been reported, they have yet to be clearly and reliably linked to specific neural

circuit- or neurocognition-based impairments in HD. Some of these not-yet-replicated HD abnormalities are present in lateral prefrontal cortex, posterior cingulate gyrus, inferior temporal cortex, and several parietal lobe regions (2-5). There also is initial evidence that patients with HD have abnormally low network functional connectivity within frontoparietal cognitive control neural systems and disrupted default mode network integrity (6). Although findings have begun to converge across HD studies on cingulo-opercular regions within the brain's salience network as one clear circuit-based HD abnormality, it is not yet clear what role these other brain function abnormalities play in HD.

One question that has not yet been extensively addressed involves the possibility that some HD brain function abnormalities might reflect the emotional state of patients with HD during symptom provocation. Cognitive behavioral theories of HD posit that maladaptive beliefs about, and emotional attachment

FEEDBA

to, possessions contribute to emotional distress (7). Consistent with these theories, patients with HD report significant subjective distress when discarding personal possessions (2,8). It would be useful to learn whether any of the functional brain abnormalities identified in prior functional magnetic resonance imaging (fMRI) studies reflect this hoarding-related distress. One relevant neuroimaging study conducted in participants without HD demonstrated that hoarding-relevant symptom provocation activated ventral prefrontal cortex and amygdala regions, suggesting that these regions may be specifically associated with hoarding-related distress (9). Other studies in HD and obsessive-compulsive disorder have linked subjective distress ratings (anxiety and sadness) and neural activity during hoarding symptom provocation in left precentral gyrus, right middle and inferior frontal gyri, and anterior ventromedial prefrontal cortex (2,4,10). However, these were post hoc analyses of primary study group differences that leave questions as to the full scope of possible HD distress/brain dysfunction relationships. To our knowledge, all prior studies that have shed light on relationships between hoarding-related distress and neural activity have focused on brain activation data, not functional connectivity. Given the emerging understanding of the importance of distributed brain network models to understanding psychopathology, it will be especially informative to determine how hoarding-related distress might be associated with abnormalities in functional connectivity.

The purpose of the present study was to learn whether hoarding-related emotional distress is associated with specific profiles of abnormal functional connectivity during a symptom provocation fMRI challenge. Specifically, we examined relationships between subjective distress during item discarding/acquiring decisions and functional connectivity across the entire brain in patients with HD and in healthy control (HC) participants. We reexamined fMRI data from a recent study that compared brain activation between HD participants and HCs during both discarding and acquiring decisions (3). We used a multivariate technique that combines multimodal canonical correlation analysis with joint independent component analysis (mCCA+jICA) (11) to examine the associations between functional connectivity measurements extracted from fMRI time series data and visual analog scale (VAS) ratings across five domains relevant to HD (fear/anxiety, sadness/regret, monetary value, importance/usefulness, and sentimental attachment). In contrast to less sophisticated, univariate techniques (e.g., simple linear correlation with VAS ratings), mCCA+jICA can identify and represent complex patterns of data relationships using a relatively small handful of components. This facilitates interpretation and mitigates the type I error rate control burden that usually would be insurmountable when examining such high-dimensional brain function data for these types of relationships, making it ideally suited to assess how VAS ratings and connectivity measurements systematically covary across patients with HD and HC participants.

By design, the mCCA+jICA approach will find systematic relationships between brain connectivity and VAS ratings. As such, we expected that the technique would successfully isolate meaningful associations between functional connectivity in specific brain regions and separate VAS ratings. Of these, we focused only on relationships found by mCCA+jICA that depicted abnormalities in HD, while relationships between

distress and connectivity found not to differ between HD and HC groups were of secondary interest. We also were most interested in connectivity-VAS rating associations during discarding decisions. Our prior fMRI studies have tended to observe a greater extent and severity of HD brain activity abnormalities in discarding relative to acquiring decisions, possibly because acquiring decisions may elicit less subjective distress in patients with HD. Based on prior HD research (9), we hypothesized that VAS anxiety ratings would be associated with metrics that quantified overall brain interconnectedness for the amygdala. Consistent with previous studies (2,10), we also hypothesized that sadness/regret ratings would be associated with connectivity strength in the precentral gyrus and middle and inferior frontal gyri. Such associations would build upon these prior studies by linking specific forms of distress in HD to the role each of these different regions play within distributed brain networks.

## METHODS AND MATERIALS

### Participants

Participants were 72 right-handed adults (mean age 55.0 years, 86% female) with a primary diagnosis of HD of at least moderate severity who enrolled in a waitlist-controlled trial of group cognitive behavioral therapy ([clinicaltrials.gov](https://clinicaltrials.gov) identifier NCT01956344) for HD ("HD group"). An age- and sex-equivalent HC group of 44 right-handed adults (mean age 54.0 years, 77% female) also participated. See Table 1 for participant characteristics. Participants were enrolled between October 22, 2013, and November 27, 2017, and have been previously described in other studies (3,12). The HD sample reported here differs slightly from the overall available sample examined in the report of Stevens *et al.* (3) due to VAS and fMRI data availability (see Supplement for details). The HD and HC groups differed on HD symptom severity (more severe in HD) and race (more racial diversity based on  $\chi^2$  analyses in the HC group), but not on ethnicity or other demographic variables (see Table 1). Full study inclusion/exclusion criteria are detailed in the Supplement.

### Study Procedure Overview

All study procedures were approved by the Hartford Hospital's Institutional Review Board. On the first day of the study, the informed consent form was reviewed with participants in detail, after which written informed consent was obtained. Participants then completed the intake interview. If eligible for the study, HD group participants returned to the laboratory approximately 1 week later for the MRI session, during which they completed discarding and acquiring fMRI symptom provocation decision-making tasks (see description, below). Immediately after each of these two tasks, they provided their VAS ratings. They also completed self-report measures of HD severity during this visit. The HC group completed the interview, fMRI tasks, VAS ratings, and self-report measures on the same day. An entire fMRI session took approximately 90 minutes, and each participant received \$50 for completing the session.

### Measures

**Diagnostic Assessments.** We used the Diagnostic Interview for Anxiety, Mood, and Obsessive-Compulsive and

**Table 1. Sample Characteristics**

Variable	HD Group, n = 72	HC Group, n = 44	Comparison, t or $\chi^2$ (p)
Age, Years, Mean (SD)	55.26 (8.74)	53.97 (7.12)	0.82 (.414)
Female Sex, n (%)	62 (86.1)	34 (77.3)	1.495 (.221)
Race, n (%)			10.182 (.006)
White	69 (95.8)	34 (77.3)	
Black	2 (2.8)	9 (20.5)	
Asian	1 (1.4)	1 (2.3)	
Hispanic Ethnicity, n (%)	2 (2.8)	3 (6.8)	1.081 (.298)
Comorbid Diagnoses, n (%)			
OCD	10 (13.9)		
Anxiety disorder	25 (34.7)		
Depressive disorder	39 (54.2)		
ADHD	12 (16.7)		
Psychiatric Medications, n (%)			
Antidepressants	13 (18.1)		
Benzodiazepines	7 (9.7)		
Stimulants	5 (6.9)		
SI-R Saving, Mean (SD)	20.06 (3.71)	3.59 (2.85)	25.23 (<.001)
SI-R Clutter, Mean (SD)	25.89 (5.79)	2.52 (3.17)	28.04 (<.001)
SI-R Acquiring, Mean (SD)	15.51 (5.51)	2.75 (2.16)	17.57 (<.001)
SI-R Total, Mean (SD)	61.46 (10.95)	11.36 (5.84)	32.07 (<.001)
Discarding Task VAS, Mean (SD)			
Fear/anxiety	23.71 (22.51)	2.09 (6.49)	7.65 (<.001)
Sadness/regret	15.74 (18.79)	1.16 (5.27)	6.20 (<.001)
Monetary value	31.65 (17.50)	27.77 (20.10)	1.10 (.276)
Importance/usefulness	38.03 (20.67)	26.36 (19.95)	2.99 (.003)
Sentimental value	23.06 (22.39)	11.00 (15.87)	3.12 (.002)
Acquiring Task VAS, Mean (SD)			
Fear/anxiety	17.85 (19.32)	3.64 (8.24)	5.48 (<.001)
Sadness/regret	9.81 (18.50)	0 (0)	4.50 (<.001)
Monetary value	31.99 (18.74)	25.11 (18.90)	1.99 (.049)
Importance/usefulness	38.40 (21.19)	28.11 (23.25)	2.45 (.016)
Sentimental value	15.75 (17.32)	6.14 (14.05)	3.26 (.002)

ADHD, attention-deficit/hyperactivity disorder; HC, healthy control; HD, hoarding disorder; OCD, obsessive-compulsive disorder; SI-R, Saving Inventory-Revised; VAS, visual analog scale.

Related Neuropsychiatric Disorders (13), a structured diagnostic interview based on the DSM-5, to assess participants' diagnoses. Because moderate HD severity was required for patient inclusion, we assessed global illness severity using a modified version of the Clinical Global Impression scale (14,15). Interviewers were licensed psychologists or psychology postdoctoral fellows under supervision.

**Self-report Measures.** Self-reported HD severity was assessed with the Saving Inventory-Revised (SI-R) (16), which contains three subscales that assess the primary symptoms of HD (acquiring, difficulty discarding, and clutter). Items are rated on a 5-point scale with higher scores indicating greater HD severity. The SI-R has demonstrated adequate internal consistency in previous studies (16) and showed excellent internal consistency in this sample (total score,  $\alpha = .98$ ; subscales, all  $\alpha$  values  $\geq .95$ ).

**Visual Analog Scale.** Participants were asked the following questions immediately after the discarding and acquiring fMRI

tasks, making a total of 10 VAS ratings (5 from each task) available for analysis. The questions were presented on the computer screen in the MRI scanner using a VAS of 0 to 100, where 0 = not at all and 100 = extremely:

*Fear/anxiety:* How much fear or anxiety did you experience?

*Sadness/regret:* How much sadness or regret did you experience?

*Monetary value:* How much monetary value did these items seem to have overall?

*Importance/usefulness:* How much importance or usefulness did these items seem to have overall?

*Sentimental attachment:* How much sentimental value did these items seem to have overall?

**fMRI Paradigms.** The discarding and acquiring tasks were based on the tasks described in Preston *et al.* (17). The tasks are detailed in prior reports (3,18) and in the Supplement. Briefly, in the discarding task, participants were presented with

pictures of household items and were asked to decide either to keep or to discard them as part of spring cleaning. In the acquiring task, participants saw pictures of items that could be acquired for free and were asked to decide whether or not to acquire them. In a control condition, participants viewed pictures and were asked to decide if the object was once alive (e.g., a wooden spoon) or never alive (e.g., a synthetic sponge). We have shown that these simulated acquiring and discarding decisions provoke excessive saving and acquiring behaviors in HD and are useful for probing HD-related brain dysfunction (18).

### MRI Data Collection and Preparation

Details of the Siemens Skyra 3T scanner (Erlangen, Germany) sequences and the Human Connectome Project data processing approach (19) were reserved for the study Supplement to focus here on how the functional connectivity data were generated and prepared for mCCA+jICA analyses. To quantify functional connectivity, vertices, or voxels within 387 separate brain regions were averaged for each time point in each participant's fMRI task time series. These regions included the Human Connectome Project's 360 cortical parcels (20), the Human Connectome Project's standard subcortical regions, and 11 discrete cerebellar regions that could be reliably mapped across participants from a published cerebellar parcellation (21). R software (22) was used to create Pearson cross-correlation matrices (adjacency matrices) among these brain regions for each participant's parcellated time series data. Each correlation value (i.e., edge) quantified the strength of functional connectivity between two separate regions. These edge values underwent Fisher's Z transformation and then were flattened into a single row of 74,691 edge values.

### Data Fusion With mCCA+jICA

Two-group (HD and HC), two-feature (functional connectivity and VAS ratings) mCCA+jICA (11) was conducted using the Fusion ICA Toolbox (available at <http://trendscenter.org/software/fit/>). mCCA+jICA is an approach to "fuse" different data features together. It is highly reliable, is robust to measurement noise, is more likely than several other fusion methods to uncover hidden relationships between or within modalities (23–25), and has been used successfully to discover disorder biomarkers and novel insights into disease (26–36). The methodology is described in detail in Sui *et al.* (34). The technical details of its application in the current study are described in the Supplement. Briefly, mCCA+jICA identifies jointly covarying information between two features within a multivariate context. Here we sought to discover the correspondence of different patterns of fMRI-measured functional connectivity while participants made decisions to acquire or discard items with their VAS-rated emotional state immediately after they finished all trials for each fMRI task. The focus of mCCA+jICA is on identifying how relationships between data features differ between study groups, not on finding group differences in the raw data themselves. So mCCA+jICA results are best used to link complex profiles of information to find associations that otherwise might be obscured by conventional analysis approaches, which often have insurmountable

statistical power requirements. When mCCA+jICA is executed, it typically is constrained to identify a manageable handful of associated patterns. Here, the mCCA+jICA analysis sought 8 joint components, which was judged likely to capture at least a handful of unique relationships between features given that there were only 10 VAS datapoints for each participant. After mCCA+jICA finds linked component pairs, a data back-reconstruction process expressed the contribution of each datapoint as z scores. More extreme z scores represent higher positive or negative expression within the relationship depicted in each component pair. For example, a VAS component with only one or two high z scores would be driven by participants' responses on those specific VAS questions. Furthermore, those VAS responses would be linked to whichever specific brain connectivity data features had high z scores in the other feature.

**Component Visualization.** Because the two features had different dimensionality, we aggregated mCCA+jICA results across participants in different ways to best visualize the relative importance of specific datapoints within each component pair. For the VAS ratings, it was straightforward to see which VAS questions loaded onto which components simply by depicting the sample's mean z scores. Interpretation of which VAS questions are most important to each component pair should follow commonly-used thresholds (e.g.,  $z > 2.0$ , or 2 SD above the mean). However, there were 74,691 edges in the functional connectivity adjacency matrices. So for the connectivity feature, it was useful to find a way to summarize these edge z weights in a way that focused on the overall connectivity of specific brain regions. To do this, we used a common graph theoretic algorithm in R software (centrality degree) to estimate a degree value for each of the 387 parcels for each participant. Each degree value represented that brain region's overall weighted importance to that component pair. For example, if the amygdala had numerous edges with high z weights for one component pair, but its z scores were close to 0 for another pair, the degree metric would be high in the first, and low in the second. As such, we could identify which handful of brain regions had connectivity values important to each component pair. We then evaluated a bootstrapped estimate of each degree value's probability of being at the 95th percentile of the observed distribution for each region to focus only on regions whose values were sufficiently different from its average across participants that they were unlikely to have occurred by chance. This was calculated using the OutlierDetection package in R software (37). To ensure a conservative estimate of statistical significance, the intersection of three different bootstrapping methods was used [depth (38), dispersion (39), and k-nearest neighbor (40), each calculated using a 95th percentile threshold].

**Testing mCCA+jICA Components for Group Differences.** mCCA+jICA also produces mixing coefficients for each participant. These coefficients represent how much each participant expressed the relationship depicted in each component. There is one mixing coefficient for each

**Table 2. Summary of Hypothesis Testing on mCCA + jICA Mixing Coefficients for the Discarding Task fMRI Connectivity Data and VAS Ratings**

Component Pair	Connectivity Feature					VAS Rating Feature				
	<i>r</i>	HC, <i>p</i>	HD, <i>p</i>	HC vs. HD, <i>p</i>	HD SI-R Total, <i>p</i>	HC, <i>p</i>	HD, <i>p</i>	HC vs. HD, <i>p</i>	HD SI-R Total, <i>p</i>	
1	.574 <sup>a</sup>	.0023 <sup>a</sup>	ns	.0006 <sup>a</sup>	.01	<.0001 <sup>a</sup>	.0016 <sup>a</sup>	.0001 <sup>a</sup>	.0001 <sup>a</sup>	
2	.569 <sup>a</sup>	ns	<.0001 <sup>a</sup>	.0001 <sup>a</sup>	.0352	.0483	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>	.0382	
3	.543 <sup>a</sup>	ns	.0247	.0377	.0278	ns	.0043 <sup>a</sup>	.0177	ns	
4	.466 <sup>a</sup>	<.0001 <sup>a</sup>	ns	<.0001 <sup>a</sup>	.0326	.013	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>	.0007 <sup>a</sup>	
5	.434 <sup>a</sup>	<.0001 <sup>a</sup>	ns	.0059 <sup>a</sup>	.015	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>	.0003 <sup>a</sup>	ns	
6	.420 <sup>a</sup>	.0045 <sup>a</sup>	ns	.0106	ns	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>	.0033 <sup>a</sup>	ns	
7	.349 <sup>a</sup>	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>	.0238	ns	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>	.0132	ns	
8	.260 <sup>a</sup>	ns	.0073	ns	ns	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>	ns	ns	

Component pairs are sorted in descending order of the Pearson *r* correlation strength between connectivity and VAS features. Subsequent columns list the significance *p* values of one-sample, two-sample, and post hoc correlation tests on the coefficient data.

fMRI, functional magnetic resonance imaging; HC, healthy control; HD, hoarding disorder; mCCA+jICA, multimodal canonical correlation analysis with joint independent component analysis; ns, not significant; SI-R, Saving Inventory-Revised; VAS, visual analog scale.

<sup>a</sup>Survives Bonferroni correction for multiple comparisons ( $p < .00625$ ).


participant for the VAS feature and connectivity feature for each of the 8 component pairs. These were used in two-sample *t*-test random effects analyses to test how the HD and HC groups differed. We were interested primarily in fused components where both the connectivity component and VAS rating component differed between groups. This would represent associations between HD brain network integration linked to a specific VAS rating profile that could be deemed abnormal in the HD group relative to the HC group. One-sample *t* tests also were done separately for each study group to evaluate whether the coefficients significantly differed from 0 for HD, HC, or both groups. The latter can be useful to provide additional interpretive context, as not every component or fused relationship between features necessarily will be strongly expressed in both groups. For instance, the presence of a brain connectivity profile in HC participants that is absent in HD can be straightforwardly interpreted as an HD deficiency. Lastly, we conducted Pearson correlations between SI-R total scores, functional connectivity, and VAS ratings for each component to learn if any features already determined to be abnormal in HD also were associated with the severity of HD symptomatology. We applied Bonferroni corrections for testing group differences between the 8 component pairs ( $\alpha .05/8 = .00625$ ).

**RESULTS**

Table 2 summarizes the results of all mCCA+jICA analyses of the discarding task fMRI data and VAS ratings. Pearson correlations between connectivity and VAS ratings across components ranged from  $r = .26$  to  $.57$ , which were all statistically significant after Bonferroni correction. This indicates that mCCA+jICA was successful in linking VAS ratings with connectivity features. Next, we present paired components of a priori interest that showed HD versus HC group differences in both functional connectivity and VAS ratings. Again, because our interest was on abnormal HD discarding-related VAS/connectivity results, acquiring fMRI task component pairs and details for any discarding task component pairs that did not differ between the HD and HC groups are described in the study Supplement.


**Group Difference Results**

Component pairs 1, 2, and 4 differed between HD and HC groups on both connectivity and VAS rating features. These met our criteria of a priori interest as they reflect HD versus HC group differences in functional connectivity and VAS ratings for these component pairs. Component pairs 1 and 4 depict a profile of brain connectivity that was abnormally absent in the



**Figure 1.** Discarding task component 1 functional connectivity map (left panel) and visual analog scale (VAS) standardized score (right panel).





**Figure 2.** Discarding task component 2 functional connectivity map (left panel) and visual analog scale (VAS) standardized score (right panel).

HD group, while component pair 2 characterizes HD connectivity that was abnormally expressed (i.e., not seen in the HC group). Data reanalysis to ensure that these differences were unrelated to HD versus HC racial/ethnic differences confirmed the results (see Supplement).

**Component Pair Description**

Figures 1 to 3 depict the functional connectivity fMRI data and the VAS scores for the 3 component pairs of interest. Table 3 lists the specific brain parcels in the fMRI feature that survived the bootstrapping test for their statistical significance. Figure 1 shows that component pair 1 was driven primarily by high fear/anxiety VAS scores. As seen in Table 3 and Figure 1, this component comprised connectivity in largely right hemisphere-lateralized dorsolateral, orbitofrontal, and frontopolar brain regions along with left nucleus accumbens and pallidum, lateral temporal cortex, and medial temporal cortex. Component pair 2 was characterized by high sadness/regret after the discarding task but also was linked to sentimental attachment VAS ratings after completing the acquiring task. This profile of VAS ratings was linked to connectivity of the bilateral amygdala, left accumbens and pallidum, bilateral medial temporal lobe, and bilateral dorsolateral prefrontal cortex (Table 3 and Figure 2). Finally, component pair 4 comprised high VAS ratings about the importance/usefulness of items discarded and connectivity strength in several lateral


prefrontal cortex and medial and lateral temporal cortex parcels (Table 3 and Figure 3).

**Correlations With Hoarding Symptom Severity**

Table 2 also reports Pearson correlations between SI-R total scores and connectivity features and VAS ratings for the discarding task fMRI data. SI-R scores were significantly correlated with the VAS loading scores for all three of these component pairs, surviving corrections for multiple comparisons for component pairs 1 and 4. There also were significant associations between SI-R scores and the brain connectivity features for all three component pairs on which we focused (component pairs 1, 2, and 4), but none of these correlations survived Bonferroni correction. These results are evidence not only that the subjective distress features here predict self-reported HD severity but also that their associated abnormal HD connectivity profiles do as well, albeit not as robustly.

**DISCUSSION**

The present study revealed several abnormal brain activity profiles that were linked to different types of subjective distress during discarding decisions in HD. Fear/anxiety VAS ratings were associated with connectivity in largely right hemisphere-lateralized dorsolateral, orbitofrontal, and frontopolar brain regions along with left nucleus accumbens and



**Figure 3.** Discarding task component 4 functional connectivity map (left panel) and visual analog scale (VAS) standardized score (right panel).

**Table 3. List of Human Connectome Project Brain Parcels That Were Identified By Bootstrapping Analysis as Having Significantly Strong Connectivity Representation for Each Pair of Fused Components From the mCCA + jICA Analysis**

Component	Hemisphere	Abbreviation	Name	Region	Super-region	Weighted Degree
Discard 1	Left	-	Pallidum	Subcortical	Subcortical	673
Discard 1	Left	PI	Parainsular area	Insular and frontal opercular cortex	Insula/frontal operculum	581
Discard 1	Left	pOFC	Posterior OFC complex	Orbital and polar frontal cortex	Frontal	557
Discard 1	Right	TGv	Area TG ventral	Lateral temporal cortex	Temporal	548
Discard 1	Right	10d	Area 10d	Orbital and polar frontal cortex	Frontal	510
Discard 1	Left	-	Accumbens	Subcortical	Subcortical	505
Discard 1	Right	TE2a	Area TE2 anterior	Lateral temporal cortex	Temporal	494
Discard 1	Right	9a	Area 9 anterior	Dorsolateral prefrontal cortex	Frontal	489
Discard 1	Right	a10p	Area anterior 10p	Orbital and polar frontal cortex	Frontal	480
Discard 1	Right	p10p	Area posterior 10p	Orbital and polar frontal cortex	Frontal	479
Discard 1	Right	8Av	Area 8Av	Dorsolateral prefrontal cortex	Frontal	478
Discard 1	Right	PeEc	Perirhinal ectorhinal cortex	Medial temporal cortex	Temporal	477
Discard 1	Right	8C	Area 8C	Dorsolateral prefrontal cortex	Frontal	469
Discard 1	Right	9p	Area 9 posterior	Dorsolateral prefrontal cortex	Frontal	460
Discard 1	Left	p10p	Area posterior 10p	Orbital and polar frontal cortex	Frontal	459
Discard 1	Left	10d	Area 10d	Orbital and polar frontal cortex	Frontal	457
Discard 2	Right	TGv	Area TG ventral	Lateral temporal cortex	Temporal	830
Discard 2	Left	-	Pallidum	Subcortical	Subcortical	636
Discard 2	Left	PI	Parainsular area	Insular and frontal opercular cortex	Insula/frontal operculum	583
Discard 2	Right	9a	Area 9 anterior	Dorsolateral prefrontal cortex	Frontal	567
Discard 2	Right	PeEc	Perirhinal ectorhinal cortex	Medial temporal cortex	Temporal	549
Discard 2	Right	EC	Entorhinal cortex	Medial temporal cortex	Temporal	535
Discard 2	Left	-	Accumbens	Subcortical	Subcortical	532
Discard 2	Left	PeEc	Perirhinal ectorhinal cortex	Medial temporal cortex	Temporal	526
Discard 2	Right	TGd	Area TG dorsal	Lateral temporal cortex	Temporal	507
Discard 2	Right	-	Amygdala	Subcortical	Subcortical	488
Discard 2	Right	TE2a	Area TE2 anterior	Lateral temporal cortex	Temporal	475
Discard 2	Right	V6A	Area V6A	Dorsal stream visual cortex	Visual	471
Discard 2	Left	TGd	Area TG dorsal	Lateral temporal cortex	Temporal	466
Discard 2	Left	9p	Area 9 posterior	Dorsolateral prefrontal cortex	Frontal	457
Discard 2	Right	9p	Area 9 posterior	Dorsolateral prefrontal cortex	Frontal	457
Discard 2	Left	-	Amygdala	Subcortical	Subcortical	448
Discard 2	Left	8Av	Area 8Av	Dorsolateral prefrontal cortex	Frontal	437
Discard 4	Right	V6A	Area V6A	Dorsal stream visual cortex	Visual	631
Discard 4	Right	TGv	Area TG ventral	Lateral temporal cortex	Temporal	611
Discard 4	Right	-	Cerebellum 3	Cerebellum	Cerebellum	589
Discard 4	Right	EC	Entorhinal cortex	Medial temporal cortex	Temporal	580
Discard 4	Right	TGd	Area TG dorsal	Lateral temporal cortex	Temporal	544
Discard 4	Left	7PI	Lateral area 7P	Superior parietal cortex	Parietal	513
Discard 4	Right	TE2p	Area TE2 posterior	Lateral temporal cortex	Temporal	500
Discard 4	Right	9p	Area 9 posterior	Dorsolateral prefrontal cortex	Frontal	480
Discard 4	Right	46	Area 46	Dorsolateral prefrontal cortex	Frontal	480
Discard 4	Right	IFSp	Area IFSp	Inferior frontal cortex	Frontal	477
Discard 4	Left	TE2p	Area TE2 posterior	Lateral temporal cortex	Temporal	462
Discard 4	Left	STSvp	Area STSv posterior	Auditory association cortex	Auditory	458
Discard 4	Left	MT	Middle temporal area	MT+ and neighboring visual areas	Visual	447
Discard 4	Left	9p	Area 9 posterior	Dorsolateral prefrontal cortex	Frontal	439
Discard 4	Right	STSvp	Area STSv posterior	Auditory association cortex	Auditory	436

Each parcel is describing using a brief parcel abbreviation and name along with anatomical region and super-region categories as described in the report of Glasser *et al.* (20). The weighted degree value for each parcel is also listed.

IFS, inferior frontal sulcus; MT+, middle temporal complex; OFC, orbitofrontal cortex; STSv, ventral superior temporal sulcus; TE2, inferior temporal gyrus; TG, temporal gyrus.

pallidum, lateral temporal cortex, and medial temporal cortex. Importance/usefulness ratings were associated with connectivity strength in similar regions. However, it should be noted that while the specific VAS rating profiles for fear/anxiety and importance/usefulness were found in both the HC and HD groups, the connectivity features associated with these ratings were found only in HC participants. Thus, participants with HD appear to lack patterns of normal connectivity. Based on the regions found in these two component pairs, one can speculate that patients with HD might not have normal interregional communication in many different functional contexts related to emotional reaction and expression [e.g., fear generalization or stimulus discrimination (amygdala and medial temporal lobe)] (41–43), which should be explored in future studies. It should be noted that these frontal regions, particularly dorsolateral prefrontal cortex, have been implicated in cognitive control/emotion regulation processes [for a meta-analysis, see (44)], suggesting that patients with HD may have emotion regulation deficits, particularly during discarding decisions. Previous studies have indeed reported greater self-reported emotion dysregulation in HD groups relative to that in control groups (45–47).

Sadness, regret, and sentimental attachment ratings during discarding were associated with connectivity of the bilateral amygdala, left accumbens and pallidum, bilateral medial temporal lobe, and bilateral dorsolateral prefrontal cortex. Some of these brain regions match those known to be engaged by sadness in prior studies (48). More interestingly, they closely resemble the medial amygdala network delineated by Bickart *et al.* (49) as brain regions having strong functional connectivity to the central and medial nuclei of the amygdala. Indeed, most of the key nodes in the amygdala subnetwork overlap with parcels identified in this component pair feature (e.g., bilateral amygdala, nucleus accumbens). The anatomical connections of this centromedial amygdala subregion position it as a major output node of the limbic system (50), with strong interconnectivity to ventromedial striatum and medial temporal lobe regions (51). Interestingly, HC participants did not express this sadness–centromedial amygdala subregion relationship at all, suggesting that it is an abnormal emotional expression–network connectivity association seen in patients with HD. Alternatively, because HC participants generally expressed very low levels of sadness when discarding, another possible explanation is that this is a normal relationship that is irrelevant to HC participants in this fMRI task context. Given the absence of these associations in the HC group, these findings may again point to potential emotion regulation deficits in HD.

We were surprised that the ACC did not appear in any component. The ACC has been identified several times in prior HD neuroimaging research, emerging as perhaps the most important area of interest in our own previous fMRI studies (2,3). Since the salience network is reliably engaged in this task (3) and when detecting the emotional significance of stimuli and generating emotional responses, we would have expected to see ACC brain function linked to HD subjective distress. But these results indicate that the cingulate's connectivity with other regions is unrelated to the emotional reactions of patients with HD or any automatic attempts to regulate their feelings when discarding possessions.

### Limitations

Limitations of the current study include the use of a sample with limited racial/ethnic diversity, making it challenging to generalize the findings to individuals from minority groups. Second, participants rated their negative emotions after completing all of the discarding and acquiring tasks, not after each individual item. More frequent ratings might allow for a more nuanced understanding of changes in negative affect when making decisions about specific types of items (e.g., items of greater vs. lesser value). Third, participants were shown a standardized set of stimuli (not an individualized set based on specific hoarding concerns). It is possible that some items were irrelevant to individual participants and therefore easier to discard. Fourth, we did not include a clinical control group, so we cannot be certain that the functional connectivity deficits we observed in this study are specific to patients with HD. Fifth, our power analysis indicated that we were adequately powered to detect medium to large effects with our sample size and Bonferroni corrections. Thus, we may have been unable to detect potentially interesting effects of smaller magnitudes. Sixth, we did not control for medication use in the analyses. Only a minority of HD group participants (22%) were taking psychiatric medications, so there were insufficient numbers of any single medication to make a statistical control approach viable. We also did not control for comorbid diagnoses in the analyses. Last, we did not use an analysis method that was able to differentiate connectivity related to the control task within the fMRI paradigms we used. So it remains possible that some of the functional connectivity–VAS findings might be not be specific to discarding or acquiring. While it seems unlikely that the control task would significantly influence the affective state of patients with HD, future research should keep the possibility in reserve.

### ACKNOWLEDGMENTS AND DISCLOSURES

This work was supported by the National Institute of Mental Health (Grant No. R01 MH101163 [to DFT and MCS]).

We thank Katrina Abertz, Amber Billingsley, Akanksha Das, Marla Genova, Benjamin Katz, Krishna Pancholi, James Ransom, and Kathryn Young for their assistance with data collection; Drs. Elizabeth Davis, Christina Gilliam, Scott Hannan, Kristen Springer, and Blaise Worden for serving as study therapists or independent evaluators; and Drs. Jonathan Abramowitz, John Goethe, and Vincent Calhoun for serving as the data and safety monitoring board.

The authors report no biomedical financial interests or potential conflict of interest.

### ARTICLE INFORMATION

From the Anxiety Disorders Center (HCL, JH, BMW, LSH, DFT) and Olin Neuropsychiatry Research Center (AP, MCS), Institute of Living, Hartford; and Department of Psychiatry (DFT, MCS), Yale School of Medicine, New Haven, Connecticut; Discipline of Clinical Psychology (BMW), Graduate School of Health, University of Technology Sydney, Sydney, South Wales, Australia; and Department of Psychology (LSH), University of Pittsburgh, Pittsburgh, Pennsylvania.

Address correspondence to Hannah C. Levy, Ph.D., at [hannah.levy@ihchealth.org](mailto:hannah.levy@ihchealth.org).

Received Oct 13, 2020; revised Dec 17, 2020; accepted Dec 23, 2020. Supplementary material cited in this article is available online at <https://doi.org/10.1016/j.bpsc.2020.12.014>.



## REFERENCES

- American Psychiatric Association (2013): Diagnostic and Statistical Manual of Mental Disorders, 5th ed. Washington, DC: American Psychiatric Publishing.
- Tolin DF, Stevens MC, Villavicencio AL, Norberg MM, Calhoun VD, Frost RO, *et al.* (2012): Neural mechanisms of decision making in hoarding disorder. *Arch Gen Psychiatry* 69:832–841.
- Stevens MC, Levy HC, Hallion LS, Wootton BM, Tolin DF (2020): Functional neuroimaging test of an emerging neurobiological model of hoarding disorder. *Biol Psychiatry Cogn Neurosci Neuroimaging* 5:68–75.
- An SK, Mataix-Cols D, Lawrence NS, Wooderson S, Giampietro V, Speckens A, *et al.* (2009): To discard or not to discard: The neural basis of hoarding symptoms in obsessive-compulsive disorder. *Mol Psychiatry* 14:318–331.
- Saxena S, Brody AL, Maidment KM, Smith EC, Zohrabi N, Katz E, *et al.* (2004): Cerebral glucose metabolism in obsessive-compulsive hoarding. *Am J Psychiatry* 161:1038–1048.
- Levy HC, Stevens MC, Glahn DC, Panchoi K, Tolin DF (2019): Distinct resting state functional connectivity abnormalities in hoarding disorder and major depressive disorder. *J Psychiatr Res* 113:108–116.
- Steketee G, Frost RO, Kyrios M (2003): Cognitive aspects of compulsive hoarding. *Cognit Ther Res* 27:463–479.
- Frost RO, Ong C, Steketee G, Tolin DF (2016): Behavioral and emotional consequences of thought listing versus cognitive restructuring during discarding decisions in hoarding disorder. *Behav Res Ther* 85:13–22.
- Mataix-Cols D, Cullen S, Lange K, Zelaya F, Andrew C, Amaro E, *et al.* (2003): Neural correlates of anxiety associated with obsessive-compulsive symptom dimensions in normal volunteers. *Biol Psychiatry* 53:482–493.
- Mataix-Cols D, Wooderson S, Lawrence N, Brammer MJ, Speckens A, Phillips ML (2004): Distinct neural correlates of washing, checking, and hoarding symptom dimensions in obsessive-compulsive disorder. *Arch Gen Psychiatry* 61:564–576.
- Sui J, Adali T, Pearson G, Yang H, Sponheim SR, White T, *et al.* (2010): A CCA+ICA based model for multi-task brain imaging data fusion and its application to schizophrenia. *Neuroimage* 51:123–134.
- Tolin DF, Wootton BM, Levy HC, Hallion LS, Worden BL, Diefenbach GJ, *et al.* (2019): Efficacy and mediators of a group cognitive-behavioral therapy for hoarding disorder: A randomized trial. *J Consult Clin Psychol* 87:590–602.
- Tolin DF, Gilliam C, Wootton BM, Bowe W, Bragdon LB, Davis E, *et al.* (2018): Psychometric properties of a structured diagnostic interview for DSM-5 anxiety, mood, and obsessive-compulsive and related disorders. *Assessment* 25:3–13.
- Guy W (1976): *Assessment Manual for Psychopharmacology*. Washington, DC: U.S. Government Printing Office.
- Tolin DF, Gilliam CM, Davis E, Springer K, Levy HC, Frost RO, *et al.* (2018): Psychometric properties of the Hoarding Rating Scale-Interview. *J Obsessive Compuls Relat Disord* 16:76–80.
- Frost RO, Steketee G, Grisham J (2004): Measurement of compulsive hoarding: Saving Inventory-Revised. *Behav Res Ther* 42:1163–1182.
- Preston SD, Muroff JR, Wengrovitz SM (2009): Investigating the mechanisms of hoarding from an experimental perspective. *Depress Anxiety* 26:425–437.
- Levy HC, Stevens MC, Tolin DF (2019): Validation of a behavioral measure of acquiring and discarding in hoarding disorder. *J Psychopathol Behav Assess* 41:135–143.
- Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, *et al.* (2013): The minimal preprocessing pipelines for the Human Connectome Project. *Neuroimage* 80:105–124.
- Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, *et al.* (2016): A multi-modal parcellation of human cerebral cortex. *Nature* 536:171–178.
- Buckner RL, Krienen FM, Castellanos A, Diaz JC, Yeo BT (2011): The organization of the human cerebellum estimated by intrinsic functional connectivity. *J Neurophysiol* 106:2322–2345.
- R Core Team (2013): *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Calhoun VD, Liu J, Adali T (2009): A review of group ICA for fMRI data and ICA for joint inference of imaging, genetic, and ERP data. *Neuroimage* 45:S163–S172.
- Calhoun VD, Sui J (2016): Multimodal fusion of brain imaging data: A key to finding the missing link(s) in complex mental illness. *Biol Psychiatry Cogn Neurosci Neuroimaging* 1:230–244.
- Sui J, Adali T, Yu Q, Chen J, Calhoun VD (2012): A review of multi-variate methods for multimodal fusion of brain imaging data. *J Neurosci Methods* 204:68–81.
- Correa NM, Eichele T, Adali T, Li YO, Calhoun VD (2010): Multi-set canonical correlation analysis for the fusion of concurrent single trial ERP and functional MRI. *Neuroimage* 50:1438–1445.
- Griffis JC, Nenert R, Allendorfer JB, Szafarski JP (2017): Linking left hemispheric tissue preservation to fMRI language task activation in chronic stroke patients. *Cortex* 96:1–18.
- Harms MP, Wang L, Csemansky JG, Barch DM (2013): Structure-function relationship of working memory activity with hippocampal and prefrontal cortex volumes. *Brain Struct Funct* 218:173–186.
- Kim SG, Jung WH, Kim SN, Jang JH, Kwon JS (2015): Alterations of gray and white matter networks in patients with obsessive-compulsive disorder: A multimodal fusion analysis of structural MRI and DTI using mCCA+jICA. *PLoS One* 10:e0127118.
- Lerman-Sinkoff DB, Sui J, Rachakonda S, Kandala S, Calhoun VD, Barch DM (2017): Multimodal neural correlates of cognitive control in the Human Connectome Project. *Neuroimage* 163:41–54.
- Mangalathu-Arumana J, Lieberthal E, Beardsley SA (2018): Optimizing within-subject experimental designs for jICA of multi-channel ERP and fMRI. *Front Neurosci* 12:13.
- Ouyang X, Chen K, Yao L, Hu B, Wu X, Ye Q, *et al.* (2015): Simultaneous changes in gray matter volume and white matter fractional anisotropy in Alzheimer's disease revealed by multimodal CCA and joint ICA. *Neuroscience* 301:553–562.
- Soh P, Narayanan B, Khadka S, Calhoun VD, Keshavan MS, Tamminga CA, *et al.* (2015): Joint coupling of awake EEG frequency activity and MRI gray matter volumes in the psychosis dimension: A BSNIP study. *Front Psychiatry* 6:182.
- Sui J, Pearson G, Caprihan A, Adali T, Kiehl KA, Liu J, *et al.* (2011): Discriminating schizophrenia and bipolar disorder by fusing fMRI and DTI in a multimodal CCA+ joint ICA model. *Neuroimage* 57:839–855.
- Sui J, Pearson GD, Du Y, Yu Q, Jones TR, Chen J, *et al.* (2015): In search of multimodal neuroimaging biomarkers of cognitive deficits in schizophrenia. *Biol Psychiatry* 78:794–804.
- Sui J, Qi S, van Erp TGM, Bustillo J, Jiang R, Lin D, *et al.* (2018): Multimodal neuromarkers in schizophrenia via cognition-guided MRI fusion. *Nat Commun* 9:3028.
- Tiwari V, Kashikar A (2019): OutlierDetection: Outlier Detection R package version 0.1.1.
- Johnson T, Kwok I, Ng R (1998): *Fast Computation of 2D Dimensional Depth Contours*. New York: Proc Int Conf on Knowledge Discovery and Data Mining (KDD), 224–228.
- Jin W, Tung A, Han J (2001): Mining top-n local outliers in large databases. San Francisco, CA: ACM SIGKDD Int Conf on Knowledge Discovery and Data Mining, 293–298.
- Hautamaki V, Karkkainen I, Franti P (2004): Outlier detection using k-nearest neighbour graph. Proc IEEE Int Conf on Pattern Recognition (ICPR). Cambridge, UK.
- LeDoux JE (2012): Evolution of human emotion: A view through fear. *Prog Brain Res* 195:431–442.
- Corodimas KP, LeDoux JE (1995): Disruptive effects of posttraining perirhinal cortex lesions on conditioned fear: Contributions of contextual cues. *Behav Neurosci* 109:613–619.
- Sangha S, Diehl MM, Bergstrom HC, Drew MR (2020): Know safety, no fear. *Neurosci Biobehav Rev* 108:218–230.

44. Buhle JT, Silvers JA, Wager TD, Lopez R, Onyemekwu C, Kober H, *et al.* (2014): Cognitive reappraisal of emotion: A meta-analysis of human neuroimaging studies. *Cereb Cortex* 24:2981–2990.
45. Fernandez de la Cruz L, Landau D, Iervolino AC, Santo S, Pertusa A, Singh S, *et al.* (2013): Experiential avoidance and emotion regulation difficulties in hoarding disorder. *J Anxiety Disord* 27:204–209.
46. Tolin DF, Levy HC, Wootton BM, Hallion LS, Stevens MC (2018): Hoarding disorder and difficulties in emotion regulation. *J Obsessive Compuls Relat Disord* 16:98–103.
47. Worden B, Levy HC, Das A, Katz BW, Stevens M, Tolin D (2019): Perceived emotion regulation and distress tolerance in patients with hoarding disorder. *J Obsessive Compuls Relat Disord* 22:100441.
48. Arias JA, Williams C, Raghvani R, Aghajani M, Baez S, Belzung C, *et al.* (2020): The neuroscience of sadness: A multidisciplinary synthesis and collaborative review. *Neurosci Biobehav Rev* 111:199–228.
49. Bickart KC, Hollenbeck MC, Barrett LF, Dickerson BC (2012): Intrinsic amygdala-cortical functional connectivity predicts social network size in humans. *J Neurosci* 32:14729–14741.
50. Pape HC, Pare D (2010): Plastic synaptic networks of the amygdala for the acquisition, expression, and extinction of conditioned fear. *Physiol Rev* 90:419–463.
51. Roy AK, Shehzad Z, Margulies DS, Kelly AM, Uddin LQ, Gotimer K, *et al.* (2009): Functional connectivity of the human amygdala using resting state fMRI. *Neuroimage* 45:614–626.