CORRECTION Open Access



Correction: Experiences of antenatal care practices to reduce stillbirth: surveys of women and healthcare professionals prepost implementation of the Safer Baby Bundle

Christine Andrews^{1*}, Frances M. Boyle^{1,2}, Ashley Pade¹, Philippa Middleton^{1,3,12}, David Ellwood^{1,4,13}, Adrienne Gordon^{1,5}, Miranda Davies-Tuck^{1,6,7}, Caroline Homer^{1,8}, Alison Griffin⁹, Michael Nicholl^{5,10}, Kirstine Sketcher-Baker¹¹ and Vicki Flenady¹

Correction: BMC Pregnancy Childbirth 24, 520 (2024) https://doi.org/10.1186/s12884-024-06712-8

Following publication of the original article [1], the authors reported an error in Fig. 2. The colours in the key representing the pre/post bars in the graph for Fig. 2. are

incorrect (around the wrong way). This relates to the pre/post-intervention findings and it is very important for this to be corrected.

The online version of the original article can be found at https://doi.org/10.1186/s12884-024-06712-8.

*Correspondence:

Christine Andrews

christine.andrews@mater.uq.edu.au

¹Centre of Research Excellence in Stillbirth, Mater Research Institute, The University of Queensland, Brisbane, Australia

²Institute for Social Science Research, The University of Queensland, Brisbane, Australia

³South Australian Health and Medical Research Institute, Adelaide,

⁴School of Medicine and Dentistry, Griffith University, Queensland,

⁵School of Medicine, The University of Sydney, Sydney, Australia

⁶The Ritchie Centre, Hudson Institute of Medical Research, Melbourne,

⁷Department of Obstetrics and Gynaecology, Monash University, Melbourne, Australia

⁸Burnet Institute, Melbourne, Australia

⁹QIMR Berghofer Medical Research Institute, Brisbane, Australia

¹⁰Clinical Excellence Commission, NSW Health, Sydney, Australia

¹¹Clinical Excellence Queensland, Brisbane, Australia

¹²The University of Adelaide, Adelaide, Australia

¹³Gold Coast University Hospital, Gold Coast, QLD, Australia



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

The correct figure is given below.

Healthcare Professionals




Fig. 2 Conversations about stillbirth and risk reduction pre/post- SBB implementation. *p < 0.001

The original article has been corrected.

implementation of the Safer Baby Bundle. BMC Pregnancy Childbirth. 2024;24:520. https://doi.org/10.1186/s12884-024-06712-8.

Published online: 28 September 2024

References

 Andrews C, Boyle FM, Pade A, et al. Experiences of antenatal care practices to reduce stillbirth: surveys of women and healthcare professionals pre-post

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.