

*This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.*



ISSN:2161-0525

## **Journal of Environmental & Analytical Toxicology**

The International Open Access  
Journal of Environmental & Analytical Toxicology

**Special Issue Title: Toxicology of Pesticides**

### **Handling Editors**

**Francisco Sanchez Bayo**  
Australia

**Richard ORTEGA**  
France

**Available online at:** OMICS Publishing Group ([www.omicsonline.org](http://www.omicsonline.org))

This article was originally published in a journal published by OMICS Publishing Group, and the attached copy is provided by OMICS Publishing Group for the author's benefit and for the benefit of the author's institution, for commercial/research/educational use including without limitation use in instruction at your institution, sending it to specific colleagues that you know, and providing a copy to your institution's administrator.

All other uses, reproduction and distribution, including without limitation commercial reprints, selling or licensing copies or access, or posting on open internet sites, your personal or institution's website or repository, are requested to cite properly.

Digital Object Identifier: <http://dx.doi.org/10.4172/2161-0525.S4-001>

# Time-Dependent Toxicity of Neonicotinoids and Other Toxicants: Implications for a New Approach to Risk Assessment

Henk A. Tennekes\* and Francisco Sánchez-Bayo

Experimental Toxicology Services (ETS) Nederland BV, 7201 KN, Zutphen, The Netherlands, and Centre for Ecotoxicology, University of Technology Sydney, Lidcombe, NSW 4141, Australia

## Introduction

A fundamental goal of toxicology is to determine safe levels of exposure to potentially poisonous substances for humans and the environment [1]. Traditionally, safe levels have been estimated in laboratory toxicity bioassays by calculating the non-observable effect level (NOEL) of a chemical to a variety of organisms which are representative of certain taxa, i.e. mammals, birds, fish, crustaceans, algae, etc. There are, however, fundamental problems with the validity of this approach, both conceptual and statistical in nature, as indicated by Landis and Chapman [2] and other authors [3]. Thus, the outdated NOEL concept is being replaced by the no-effect concentration (NEC) level [4], which assumes that toxic chemicals do not have any effect on a population of organisms at very low concentrations.

Druckrey and Küpfmüller [5] reasoned that the following processes determine the complex relationship between exposure levels to poisons and a toxic effect:

1. **Absorption, distribution, metabolism and excretion**, which determine the relationship between dose and the concentration of the poison at the site of action and its time course in various parts of the body, commonly referred to as pharmacokinetics. A linear relationship between dose and effect would require proportionality between dose (or concentration in a medium) and the concentration of the poison at the site of action. This may be the case for many poisons, for as long as the poison does not influence pharmacokinetics within the considered dose range;
2. **The interaction of the poison with critical receptors**, which determines the relationship between the concentration of the poison at the site of action and receptor binding. A linear relationship between dose and effect would require proportionality between the concentration of the poison at the site of action and the concentration of bound receptors. This is only the case, as will be shown in section 2, when the proportion of bound receptors sufficient for an effect is relatively small compared to the initial level of the receptor; and
3. **Receptor binding**, which determines the relationship between the concentration of bound receptors and the actual biological effect. A linear relationship between dose and effect would require proportionality between relative receptor binding and the actual biological effect.

The pharmacokinetics of most pesticides is well known after having been the subject of intense research in recent decades. For example, the kinetics of acetylcholinesterase inhibitors such as carbamate and organophosphorus insecticides was described long ago, and can be found in most toxicology textbooks (e.g. [6]). The biochemical interactions between these insecticides and their target enzyme (acetylcholinesterase), together with the kinetics of uptake, form the basis of mechanistic models of toxicity such as DEBtox [7]. In a similar way, the kinetics of narcotic chemicals has been described by models which consider the time course in their uptake and bioconcentration by fish [8].

In this review we will focus on the relationship between the concentration of the poison at the site of action and the actual biological effect, which Druckrey has referred to as *ergokinetics* (Druckrey, personal communication with Tennekes, 1985). The reason for this is that recent developments in ecotoxicology suggest that some toxicants can produce effects at any concentration level provided their exposure time is sufficiently long [9]. This means that the concept of NEC may not apply for these toxicants when the life span of the organisms affected is longer than the theoretical maximum exposure time. Consequently, risk assessment of these chemicals, which includes neonicotinoid insecticides and certain metallic compounds, may require entirely new approaches.

## Conceptual model for toxicant effects with time

In their book *Dosis und Wirkung* (dose and response) [5], Druckrey and Küpfmüller analyzed ergokinetics in mathematical terms, and their reasoning leads to fundamentally important conclusions on the effects caused by interaction of poisons with critical receptors, which are highly relevant for toxicological risk assessment.

Denoting the initial concentration of critical receptors that a poison reacts with as  $R_0$ , the concentration of receptors that a poison has reacted with as  $C_R$ , and the poison concentration at the site of action as  $C$ , the velocity of receptor binding (association) is:

$$K C (R_0 - C_R) \quad (1)$$

where  $K$  is the reaction constant for association. The velocity of dissociation of bound receptors is:

$$C_R / T_R \quad (2)$$

where  $T_R$  is the time constant for dissociation. Therefore, the reaction kinetics of receptor binding in the case of a bimolecular reaction are:

$$dC_R / dt = K C (R_0 - C_R) - C_R / T_R \quad (3)$$

If the effect occurs under circumstances where  $C_R \ll R_0$ , i.e., with first order kinetics, then  $R$  remains practically constant, in which case

$$K (R_0 - C_R) = 1 / T_A \quad (4)$$

\*Corresponding author: Henk A. Tennekes, Experimental Toxicology Services (ETS) Nederland BV, Frankensteeg 4, 7201 KN, Zutphen, The Netherlands, Tel: +31 575 545 500; Fax: +31 575 516 717; E-mail: [info@toxicology.nl](mailto:info@toxicology.nl)

Received November 14, 2011; Accepted December 05, 2011; Published December 07, 2011

Citation: Tennekes HA, Sánchez-Bayo F (2011) Time-Dependent Toxicity of Neonicotinoids and Other Toxicants: Implications for a New Approach to Risk Assessment. J Environment Analytic Toxicol S4:001. doi:10.4172/2161-0525.S4-001

Copyright: © 2011 Tennekes HA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

where  $T_A$  can be regarded as the time constant for association. Equation (3) then simplifies to

$$dC_R/dt = C/T_A - C_R/T_R \quad (5)$$

In equilibrium, where  $dC_R/dt = 0$ , Equation (5) simplifies to

$$C_R = C (T_R / T_A) \quad (6)$$

Replacing the concentration of bound receptors  $C_R$  by the relative concentration of bound receptors  $C_R/R_0$ , we obtain

$$C_R / R_0 = (C / R_0) \cdot (T_R / T_A) \quad (7)$$

This expression indicates that, with first order kinetics, i.e. when  $C_R \ll R_0$ , the proportion of bound receptors is dependent on the concentration of poison in the body of the organism (exposure dose) and the relative time for association and dissociation (exposure duration). Based on this reasoning, Druckrey and Küpfmüller drew the following conclusions for all poisons that interact with specific receptors in a first order bimolecular reaction where the toxic effect is determined by the relative concentration of bound receptors  $C_R/R_0$ :

- The effect is proportional to the concentration of the poison at the site of action C (Paracelsus).
- The effect is inversely proportional to  $R_0$ . If the concentration of specific receptors  $R_0$  is low, the poison may induce pronounced toxicity at very low concentrations at the site of action C.
- The effect is proportional to  $T_R/T_A$ , i.e., to the quotient of the time constant for dissociation  $T_R$  and the time constant for association  $T_A$ .
- If  $T_R/T_A$  is high, the poison may induce pronounced toxicity at very low concentrations at the site of action C.
- If  $T_R/T_A < 1$ , a compound is innocuous and can only induce toxicity at relatively high concentrations at the site of action C.
- If both time constants ( $T_R$  and  $T_A$ ) are low, i.e. when both association and dissociation are fast processes, equilibrium between C and receptor binding (and effect) will be established quickly but the toxic effect will also regress quickly.
- The character of a poison is primarily determined by  $T_R$ , i.e., by the time constant for dissociation of bound receptors.
- If  $T_R$  is low, i.e. when receptor binding is quickly reversible, the time course of the effect will be the same as the time course of C and the maximum effect will occur when C is at its maximum. Under these circumstances, there will be a linear relationship between effect and the actual concentration of the poison at the site of action C, and between dose and effect, provided C is proportional to dose (or concentration in a medium). Such poisons can therefore be termed “concentration poisons”. The value of  $T_A$  will only determine the fraction of the poison that reacts with the specific receptors R.
- If the time constant for dissociation  $T_R$  is high, i.e. when receptor binding is only slowly reversible, the time to maximum effect will be delayed, and the (toxic) effect will also be slowly reversible. The higher is  $T_R$  is, the longer is the time to maximum effect. Upon repeated exposure in quick succession there will be cumulative effects. Such poisons can therefore be termed “cumulative poisons”. With cumulative poisons, a quantitative or even linear relationship between effect and the actual concentration of the poison at the site of action C, as seen with concentration poisons, will not exist. Because equilibrium between C and receptor binding (and effect)

will be established very slowly, toxicity becomes a process that takes place in time. There will be a latency period up to a defined effect, which can be shortened, of course, by increasing the concentration of the poison at the site of action C.

Thus, according to Druckrey and Küpfmüller, depending on the time constant for dissociation  $T_R$ , the toxicity of a compound may become a process that takes place in time. The traditional approach to toxicity testing is to consider dose (concentration)-effect relationships at arbitrarily fixed exposure durations which are supposed to reflect ‘acute’ or ‘chronic’ time scales. This approach measures the proportion of all exposed individuals responding by the end of specified exposure times. Toxicological databases established in this way are collections of endpoint values obtained at fixed times of exposure. As such these values cannot be linked to make predictions for the wide range of exposures encountered by humans or in the environment. Thus, current toxicological risk assessment can be compromised by this approach to toxicity testing, as will be demonstrated in this paper, leading to serious underestimates of actual risk.

### Time effects of toxicants

In order to overcome this handicap, an increasing number of researchers are using a variant of the traditional toxicity testing protocol which includes time to event (TTE) methods. This TTE approach measures the times to respond for all individuals, and provides information on the acquired doses as well as the exposure times needed for a toxic compound to produce any level of effect on the organisms tested [10]. Consequently, extrapolations and predictions of toxic effects for any combination of concentration and time are now made possible [11]. We will demonstrate that this approach is superior to current toxicological testing procedures, and has important implications for risk assessment of chemicals.

Following the conceptual model by Druckrey and Küpfmüller, if receptor binding is virtually irreversible, then  $T_R \rightarrow \infty$  and Eq. (3) reduces to

$$dC_R/dt = K C (R_0 - C_R) \quad (8)$$

If the toxic effect occurs when  $C_R \ll R_0$ , then

$$dC_R/dt = K R_0 C \quad (9)$$

If a dose level is kept constant throughout a study, and, as a result, C remains constant as well, integration yields

$$C_R = K R_0 C t \quad (10)$$

This reasoning by Druckrey and Küpfmüller provided a theoretical explanation for Haber’s rule [12]. Haber’s rule (for a review, see [13]) states that the product of exposure concentration and exposure duration produces a constant toxic effect. Haber had noted that exposure to a low concentration of a poisonous gas for a long time often had the same effect (death) as exposure to a high concentration for a short time. The results of Druckrey’s ground-breaking study on the carcinogenicity of 4-dimethylaminoazobenzene (4-DAB) in BDIII rats [14] were consistent with Haber’s rule: doubling the daily 4-DAB dose, and thereby presumably doubling the concentration of the carcinogen at the site of action, halved the time up to the appearance of liver cancer.

In the 1960s the molecular biology of this carcinogenic effect was elucidated and confirmed the theorem of Druckrey and Küpfmüller. DNA was recognized by Brookes and Lawley [15] as the target for chemical carcinogens, as recently inferred by Wunderlich [16], and

Warwick and Roberts confirmed irreversible receptor binding by demonstrating covalent binding of a 4-DAB metabolite to DNA [17].

It is now apparent that Haber’s rule is highly relevant for ecotoxicological risk assessment as well. It was recently shown to describe the toxicity of the neonicotinoid insecticide imidacloprid to midges *Chironomus tentans* [18]. The product of exposure concentration and exposure duration to 50% mortality (t50) for *C. tentans* was very similar under acute and chronic exposure conditions. These observations have also confirmed the theorem of Druckrey and Küpfmüller and Haber’s rule. Abbink certified in 1991, when the compound was first introduced, that “imidacloprid is the first highly effective insecticide whose mode of action has been found to derive from almost complete and virtually irreversible blockage of postsynaptic nicotinic acetylcholine receptors (nAChRs) in the central nervous system of insects” [19]. Imidacloprid mimics the action of acetylcholine, but unlike acetylcholine, imidacloprid is not deactivated by acetylcholinesterase and thus persistently activates nAChRs [20]. Chronic exposure of insects to imidacloprid therefore leads to cumulative and virtually irreversible blockage of nAChRs in their central nervous system, which play roles in many cognitive processes.

### Exposure time reinforcing the toxic effects

Although equation (10) provided a theoretical explanation for Haber’s rule, it assumed proportionality between the concentration of bound receptors  $C_R$  and the effect. This may not always be the case. As mentioned earlier, Druckrey and Küpfmüller had pointed out that the relationship between the poison concentration at the site of action  $C$  and the effect  $E$  involves at least 2 steps: the first is the interaction of the poison with critical receptors leading to receptor binding, the second is the subsequent biological effect resulting from receptor binding. As we have seen earlier, if receptor binding is virtually irreversible (i.e.  $T_R \rightarrow \infty$ ), the concentration of bound receptors  $C_R$  is proportional to the integral of  $C$  over time:

$$C_R \sim \int C dt \tag{11}$$

If the subsequent effect is irreversible as well (e.g. death), the effect  $E$  is proportional to the integral of the concentration of bound receptors  $C_R$  over time:

$$E \sim \int C_R dt \tag{12}$$

In cases of irreversible receptor binding and an irreversible effect, the effect  $E$  is thus proportional to the double integral of the poison concentration at the site of action  $C$  over time, as the combination of equation (11) and (12) shows:

$$E \sim \iint C dt \tag{13}$$

The implication is that exposure time will reinforce the effect. Reinforcement of an effect by exposure time was subsequently demonstrated by Druckrey and co-workers with benchmark studies of the production of ear duct and liver carcinomas by 4-dimethylaminostilbene (4-DAST) [21] and diethylnitrosamine (DENA) [22], respectively, in BDII rats (Table 1).

The total carcinogenic dose *decreased* with decreasing daily 4-DAST or DENA dose levels, even though the median tumor induction times *increased* with decreasing daily dose levels. In a logarithmic system of coordinates, there was a linear relationship between the median tumor induction time (t50) and the daily dosage (d):

$$\ln d = \ln k - n \ln t50 \tag{14}$$

or

$$d t50^n = \text{constant} \tag{15}$$

where the time exponent  $n$  was 3.0 and 2.3 for 4-DAST and DENA, respectively.


The essence of equation (15) is that the total dose required to produce the same effect decreases with decreasing exposure levels, even though the exposure times required to produce the same effect increase with decreasing exposure levels. So, it should be possible, in principle, to induce cancer with a single dose of a chemical carcinogen. In fact, this was achieved by Druckrey *et al* [23] in single dose experiments with the direct-acting ethylating nitrosamide N-nitroso-N-ethylurea (ENU) in BD IX rats. This carcinogen is rapidly lost from the blood after intravenous injection; it has an *in vivo* half-life of 5-6 minutes.

Upon single dose treatment in early life (1, 10 or 30 days after birth), the overwhelming majority of animals died from malignancies of the nervous system [23]. The quantitative relationship (Figure 1) between (post-conception) age at treatment (a), dose (d), and the median induction period of neurogenic malignancies (t50) can be described as follows [24]:

$$a / t50 = K d^r \tag{16}$$

Chemical	Daily dose, d (mg kg <sup>-1</sup> )	Median tumor induction time, t (days)	Total dose, D (mg kg <sup>-1</sup> )	n
4-DAST	3.4	250	850	3.0
	2.0	340	680	
	1.0	407	407	
	0.5	550	275	
	0.28	607	170	
	0.2	700	140	
	0.1	900	90	
DENA	9.6	101	963	2.3
	4.8	137	660	
	2.4	192	460	
	1.2	238	285	
	0.6	355	213	
	0.3	457	137	
	0.15	609	91	
	0.075	840	64	

**Table 1:** Induction of ear-duct carcinomas in BDII rats by 4-DAST and of liver carcinomas by DENA, after Druckrey and Dischler [21], and Druckrey *et al*. [22].



**Figure 1:** Median age-dependent velocity of malignant neurogenic tumor formation (A/T) in BD IX rats versus single dose levels of ethylnitrosourea, on logarithmic coordinates. Linearity leads to equation (17).



with  $K = \text{constant}$ , where  $r = 0.426$

or

$$d(t50/a)^n = \text{constant} \quad (17)$$

where  $n = 1/r = 2.35$ .

Equation (17) indicates that, in this case, the *velocity* of carcinogenesis is determined by *the initiating dose* and *the state of the relevant targets at treatment*. This is reminiscent of Wilder's law of initial value, which states that the direction of response of a body function to any agent depends to a large degree on the initial level of that function. Target cells for ENU in early life are subependymal cells in the brain and subpial cells in the spinal cord, which are destined to differentiate into glial cells, i.e., astrocytes or oligodendrocytes. The evidence suggests that the critical genetic changes induced by the carcinogen in these target cells lead to inhibited normal cell differentiation processes, as has been observed in leukemogenesis [25,26]. Yuspa and Morgan reported that cells resistant to terminal differentiation can be readily isolated from skin of BALB/c mice exposed to an initiating dose of carcinogen *in vivo* but not from control mouse skin [27].

## Applications of the Druckrey-Küpfmüller model

### Genotoxic carcinogens

The Druckrey-Küpfmüller equation (15) can serve as a basis for risk assessment of carcinogens. Carlborg pointed out that this equation is implied by a Weibull model for dose-response functions in carcinogenesis [28]. The simple form of the Weibull model is a sigmoid curve defined by four parameters:

$$P = 1 - e^{-\exp}, \text{ with the exponent} = -(\alpha + \beta x^m) \quad (18)$$

where  $x$  is the dose,  $P$  is the tumor rate and  $m, \alpha, \beta$  are parameters to be estimated from the data. The parameter  $\alpha$  is determined by the background tumor probability,  $\beta$  is a scale parameter related to the units measuring the dose, and  $m$  is the important shape parameter. At very low doses the excess risk over background is approximately  $\beta x^m$ . The virtual safe dose (VSD) corresponding to a one-in-a-million risk over background is then given by

$$\text{VSD} = (10^{-6} / \beta)^{1/m} \quad (19)$$

The extended form of the Weibull model includes the time to a tumor ( $t$ ):

$$P = 1 - e^{-\exp}, \text{ with the exponent} = -(\alpha + \beta x^m) t^k \quad (20)$$

where  $k$  is a new parameter. Now suppose that  $t$  measures the time to a tumor. Also, suppose that the background tumor rate is zero ( $\alpha = 0$ ). For a test group at some dose  $x$ , consider the median time to tumor  $t50$  – that is, the value of  $t$  such that  $P = 0.50$ . The extended Weibull model for this dose and time is

$$0.50 = 1 - e^{-\exp}, \text{ with the exponent} = -\beta x^m t50^k \quad (21)$$

This reduces to

$$[-\ln 0.50 / \beta]^{1/m} = x t50^{k/m} = x t50^n \quad (22)$$

where  $n = k/m$  and the left side of the equation is a constant. This is the Druckrey-Küpfmüller equation (15). The exponent  $n$  is given by the slope of the regression between the logarithms of the dose and that of the median time to tumor.

### Neonicotinoid insecticides

Similar relationships have been demonstrated for the toxicity

of neonicotinoid insecticides to aquatic invertebrates, in particular imidacloprid to the freshwater ostracod *Cypridopsis vidua* and to *Daphnia magna* and thiacloprid to *Gammarus* and *Sympetrum*. Sánchez-Bayo [29] demonstrated that the relationship between the concentration of the neonicotinoid insecticides imidacloprid and thiacloprid in a medium ( $C$ ) and the time to 50% mortality ( $t50$ ) of several species of exposed arthropods followed a hyperbolic curve described by the equation

$$t50 = a \cdot C^{-b} \quad (23)$$

Accordingly, there was a linear relationship when the logarithms of the variables  $C$  and  $t$  were used

$$\ln t50 = a' - b \ln C \quad (24)$$

where  $a'$  is the intercept and  $b$  is the slope of the regression. Equation (24) can be transformed to

$$C^b t50 = \text{constant} \quad (25)$$

or

$$C t50^{1/b} = \text{constant} \quad (26)$$

Equation (26) is very similar to the Druckrey-Küpfmüller equation (15) for the action of chemical carcinogens such as DENA or 4-DAST.

Similar to the dose-response characteristics of DENA, exposure time was found to reinforce the toxicity of imidacloprid and thiacloprid to the tested arthropod species – Beketov and Liess [30] have referred to this feature of the neonicotinoids as having “delayed effects”. The  $C t50$  product, which reflects the total dose required for a lethal effect, decreased with decreasing toxicant concentration  $C$  (Table 2), even though the times to 50% mortality  $t50$  increased with decreasing toxicant concentration  $C$ .

This toxic behaviour of neonicotinoid insecticides has also been pointed out by other researchers. For example, Suchail et al. [31] noted that at concentrations of 0.1, 1, and 10  $\mu\text{g}$  of imidacloprid per liter, the total cumulate dose ingested by honeybees in chronic intoxication was about 60 to 6,000 times lower than the doses needed to produce the same effect in acute intoxication tests. The same effect was also apparent with two of the metabolites of imidacloprid (5-hydroxyimidacloprid and olefin). Although the authors could not explain the huge discrepancies between the results from acute toxicity tests of imidacloprid and its feeding effects in honeybees, their observations are consistent with the time-dose model described here. Consequently, a correct understanding of the ergokinetics of imidacloprid is essential for its risk assessment in relation to bees. In addition, at sub-lethal doses imidacloprid can alter honey bee foraging and learning [32-35]. Imidacloprid has been detected at levels of 5.7  $\mu\text{g kg}^{-1}$  in pollen from French hives [36] and foraging honey bees reduced their visits to a syrup feeder when it was contaminated with 3  $\mu\text{g kg}^{-1}$  of imidacloprid [37]. Foraging as well as hive worker bees and brood are likely to be continuously exposed to imidacloprid when contaminated food is collected and stored inside the hive [38]. A honey bee during a foraging flight must learn and recall many complex visual patterns [39,40]. These cognitive functions may be perturbed when nAChRs, necessary for the formation of long-term memory and involved in acquisition and retrieval processes, are persistently blocked [41]. These observations are consistent with the theorem of Druckrey and Küpfmüller [5]. Both receptor binding and the effect of receptor binding are virtually irreversible, and exposure time will therefore reinforce the effect, which may in the course of time be detrimental to the bee colony, and ultimately cause colony collapse.

Species	Chemical	Concentration (C) in $\mu\text{g L}^{-1}$	Time to 50% mortality (T50) in days	C x T50 product in $\mu\text{g L}^{-1} \cdot \text{days}$	n
<i>Cypridopsis vidua</i>	Imidacloprid	4	5.2	20.8	4.67
		16	3.0	48	
		64	3.3	211.2	
		250	2.3	575	
		1,000	2.0	2,000	
		4,000	0.9	3,600	
<i>Daphnia magna</i>	Imidacloprid	250	384.7	96,175	1.35
		750	69.7	52,275	
		2,220	18.6	41,292	
		6,700	15.0	100,500	
		20,000	18.4	368,000	
		60,000	3.0	180,000	
<i>Gammarus pulex</i>	Thiacloprid*	99	63.6	6,296.4	1.11
		364	16.7	6,078.8	
		988	6.5	6,422	
		3,100	3.2	9,920	
		9,520	0.9	8,568	
<i>Sympetrum striolatum</i>	Thiacloprid*	7.2	20.6	148.3	1.53
		8.0	17.2	137.6	
		12.7	13.0	165.1	
		113.3	3.2	362.6	

\*Original data from Beketov and Liess [30]

**Table 2:** Mortality of Arthropods Induced by Neonicotinoid Insecticides (After Sánchez -Bayo [29]). Data fit the equation  $C \cdot T50^n = \text{constant}$ .

Thus, low environmental concentrations of these insecticides (that may not be acutely toxic) could be detrimental to many invertebrate species in the medium to long term, in particular to aquatic invertebrates because these compounds are quite soluble in water and their toxicity may be reinforced by exposure time. The results of recent studies are consistent with this concept. Thus, imidacloprid applied to rice mesocosms eliminated all zooplankton species for at least two months when concentrations were above  $1 \mu\text{g L}^{-1}$  [42], which is 500 to 75,000 times lower than its acute LC50s for ostracods and cladocerans. By contrast, the insecticide fipronil applied to the same mesocosms produced only small, non-significant adverse effects on zooplankton, even if this insecticide is 100–1000 times more toxic to those organisms than imidacloprid and more persistent in soil [43]. Mayflies of the genera *Baetis* and *Epeorus* showed a reduction in reproductive success when exposed to concentrations of imidacloprid as low as  $100 \text{ ng L}^{-1}$  [44]; whether this is the result of a time-exposure effect or a sublethal effect is not clear. A single pulse contamination of mesocosms with the neonicotinoid insecticide thiacloprid resulted in long-term alteration of the overall invertebrate community structure because some species did not recover [45]. One species, the stonefly *Nemoura cinerea*, was affected at the lowest tested concentration, 70 times below the lowest known LC50. Imidacloprid can be applied as a systemic insecticide to trees by direct stem injections or by soil injections and drenches, in which case it may be indirectly introduced to aquatic systems via leaf fall or leaching: at realistic concentrations in leaves ( $18\text{--}30 \mu\text{g g}^{-1}$  fresh weight) imidacloprid can inhibit leaf litter breakdown through adverse effects on decomposer invertebrates [46], but it has no effect on microbial decomposers.

Terrestrial ecosystems have also been reported as negatively affected by imidacloprid. A study conducted over 3 years on an experimental home lawn [47] revealed that three consecutive years of imidacloprid applications to the same field plots suppressed the numbers of total hexapods, *Collembola*, *Thysanoptera* and *Coleoptera* adults by 54–62%. However, applications of imidacloprid to an experimental vegetable patch had only a temporary effect on the arthropod communities of

the eggplant crop [48], perhaps because immigration of insects from nearby areas compensated the losses caused by the insecticide. When imidacloprid is applied as a systemic insecticide to the soil around trees it may cause sublethal effects on earthworms if concentrations in the litter reach or exceed  $3 \text{ mg kg}^{-1}$  [49,50]. A recent study indicates high toxicity of imidacloprid to the non-target terrestrial arthropod *Porcellio scaber*, at similar levels as the organophosphorus diazinon [51]. However, because the blocking of AChRs by diazinon is temporary (it may last only a few hours), its toxic effect would last as long as it is present in soil at sufficient concentrations, whereas imidacloprid effects could last much longer if the nAChRs are permanently blocked.

### Other chemicals

Time-dependent effects have also been observed with toxic metals. Sánchez-Bayo [29] analysed the data provided by other authors and concluded that the toxicity of copper, zinc, selenium and cadmium (as  $\text{CdCl}_2$ ) to *Daphnia magna* and that of zinc to the guppy (*Poecilia reticulata*) followed the same pattern described by equation (26). The median times to 50% mortality (t50) decreased with increasing concentrations of metals, as seen with carcinogens and neonicotinoids, suggesting that for all metals the time constant for dissociation from the critical receptor  $T_R$  is high. However, with the exception of selenium, and in complete contrast to the action of carcinogens and neonicotinoids, the total metal dose taken up by and lethal to these aquatic organisms also decreased with increasing concentrations of metals, as shown here for *Daphnia magna* (Table 3). Selenium's toxicity followed Haber's Rule, with  $n=1$ , and can safely be assumed to bind irreversibly to the critical receptor, but the n-values characterizing the toxicity of copper, zinc, and cadmium were 0.3, 0.47, and 0.60, respectively. So, whereas with carcinogens and neonicotinoids, lower concentrations are more effectively poisonous than higher concentrations, it turns out that lower metal concentrations are less effectively poisonous than higher metal concentrations. This could reflect some elimination process which minimizes metal toxicity with time of exposure.

Chemical	Concentration (C) in $\mu\text{g L}^{-1}$	Time to 50% mortality (T50) in days	C x T50 product in $\mu\text{g L}^{-1} \cdot \text{days}$	n
Copper*	28	32.6	912	0.30
	32	6.5	208	
	48	3.2	154	
	56	2.0	112	
	64	1.3	80	
CdCl <sub>2</sub> **	3.2	292	935	0.60
	5.6	58	325	
	10	38	375	
	18	11	203	
	32	6	181	
	56	2	105	
Selenium*	158	3.3	514	1.03
	800	0.7	567	
	1200	0.5	600	
	1600	0.4	600	
	2000	0.3	500	
Zinc*	59	163	9617	0.47
	125	32	4000	
	250	2	490	
	500	1.3	667	

\*Original data from Hoang et al. [64]

\*\*Original data from Kooijman [65]

**Table 3:** Mortality of *Daphnia magna* induced by metallic elements or compounds (After Sánchez -Bayo [29]). Data also fit the equation  $C \times T50^n = \text{constant}$ .

Metal toxicity is particularly insidious, as it affects not only aquatic but also most terrestrial organisms through the food chain. While more information is necessary to prove that other metallic and metalloid elements (e.g arsenic) produce time-dependent toxic effects, the evidence so far obtained from toxicity experiments carried out with fish (e.g. [52-54]) illustrates a toxicity pattern similar to that observed with neonicotinoid insecticides. This is likely to result from metal accumulation in the body, which increases with time until the organisms reach a sufficient dose to cause a toxic effect (see review in [55]). At least we know this is the case with mercury contamination in humans, which leads to the condition known as Minamata disease [56].

It is apparent that any chemical that permanently binds to a receptor in the body to produce a toxic effect may have time-dependent effects whenever the bound receptors remain in the body of the organism. The examples shown here for carcinogenic substances that bind to DNA, the neonicotinoid insecticides that deactivate the nAChRs and the metallic elements or compounds, like CdCl<sub>2</sub>, that accumulate in the organisms' tissues, suggest this is probably the case. This suggests that many other substances may behave in the same way, even though we may be unaware of their gokinetics.

So far we have discussed only toxicants that affect animals, but in principle time-dependent toxicity could also apply to the mode of action of herbicides in plants and algae. For example, if recalcitrant herbicidal compounds (e.g. PSII inhibitors) can block permanently the photosynthetic pathway in the chloroplasts, it is likely they may also show a time-dependent toxic effect in algae and higher plants. This would have serious implications for the risk assessment of herbicides in agricultural crops and coral reefs, among others. Research in this area of ecotoxicology is still lacking, and therefore we can only speculate about it.

### Implications for risk assessment

Although time-to-event models have been considered in recent

years for inclusion in risk assessments of environmental contaminants in areas as diverse as agriculture [57], occupational health [58], engineering [59] and ecology [60], the implications of the time-dependent toxicity of some chemicals have not been realised yet. First of all we need to know what kind of toxicants behave in a time-dependent manner, which is the same as asking which chemicals bind irreversibly to specific receptors. So far, the evidence pointed out here indicates that neonicotinoids, some carcinogens and metal/metalloids fit such description but other chemical groups may also follow this pattern.

Secondly, if a toxicant has time-dependent effects, the standard risk assessment procedures would not be valid in situations where there is exposure to sublethal concentrations of the toxicant for long periods of time. Traditionally this type of exposure has been considered as chronic toxicity, and its relationship with standard acute toxicity endpoints (e.g. LC50) has been studied in many aquatic species [61] ever since Kenaga introduced a ratio to describe it [62]. However, this traditional approach ignores the underlying mechanism of toxicity with time that has been described in this review.

There is no doubt that once ecotoxicologists realise the full potential and advantages of time-to-event approaches, they will become the standard tool "for analysis of toxicity from pulse exposures, and the latent toxic effects emerging after exposure has ceased, because both of these phenomena are time related" [63]. We already pointed here the experimental work by Beketov and Liess [30] in this regard, and only hope that more researchers may follow the same pathway.

### References

1. Truhaut R (1977) Ecotoxicology: objectives, principles and perspectives. *Ecotoxicol Environ Saf* 1:151-173.
2. Landis WG, Chapman PM (2011) Well past time to stop using NOELs and LOELs. *Integr Environ Assess Manage* 7: vi-viii.
3. Crane M, Newman MC (2000) What level of effect is a no observed effect? *Environ Toxicol Chem* 19: 516-519.
4. Kooijman SALM, Bedaux JJM, Slob W (1996) No-effect concentrations as a basis for ecological risk assessment. *Risk Anal* 16: 445-447.
5. Druckrey H, Küpfmüller K (1949) Dosis und Wirkung - Beiträge zur theoretischen Pharmakologie. Editio Cantor GmbH, Freiburg im Breisgau, Germany.
6. Matsumura F (1985) Toxicology of pesticides. Plenum Press, New York.
7. Kooijman SALM, Bedaux JJM (1996) The analysis of aquatic toxicity data. VU University Press, Amsterdam, The Netherlands.
8. Mackay D, Puig H, McCarty LS (1992) An equation describing the time course and variability in uptake and toxicity of narcotic chemicals to fish. *Environ Toxicol Chem* 11: 941-951.
9. Tennekes HA (2010) The significance of the Druckrey-Küpfmüller equation for risk assessment - The toxicity of neonicotinoid insecticides to arthropods is reinforced by exposure time. *Toxicology* 276: 1-4.
10. Newman MC, McCloskey JT (1996) Time-to-event analyses of ecotoxicity data. *Ecotoxicology* 5: 187-196.
11. Crane M, Newman MC, Chapman PF, Fenlon J (2002) Risk assessment with Time to Events models. CRC Press LLC, Boca Raton, FL.
12. Haber F (1924) Zur Geschichte des Gaskrieges. In: Fünf Vorträge aus den Jahren 1920-1923. Julius Springer, Berlin, pp. 76-92.
13. Witschi H (1999) Some notes on the history of Haber's law. *Toxicol Sci* 50: 164-168.
14. Druckrey H (1943) Quantitative Grundlagen der Krebszeugung. *Klin Wochenschr* 22: 532.
15. Brookes P, Lawley PD (1964) Evidence for the binding of polynuclear aromatic hydrocarbons to the nucleic acids of mouse skin: Relation between carcinogenic power of hydrocarbons and their binding to deoxyribonucleic acid. *Nature* 202: 781-784.



16. Wunderlich V (2007) "Substances capable of self-reproduction" as cellular targets of chemical carcinogens. *Ntm* 15: 271-283.
17. Warwick GP, Roberts JJ (1967) Persistent binding of Butter Yellow metabolites to rat liver DNA. *Nature* 213: 1206-1207.
18. Stoughton SJ, Liber K, Culp J, Cessna A (2008) Acute and chronic toxicity of imidacloprid to the aquatic invertebrates *Chironomus tentans* and *Hyalella azteca* under constant- and pulse-exposure conditions. *Arch Environ Contam Toxicol* 54: 662-673.
19. Abbink J (1991) The biochemistry of imidacloprid. Pflanzenschutz-Nachrichten Bayer, Germany, F.R., Serial ID – ISSN: 0340-1723
20. Thany SH (2010). Neonicotinoid insecticides. Historical evolution and resistance mechanisms. In: Thany SH (Ed.) Insect nicotinic acetylcholine receptors. *Advances in Experimental Medicine and Biology* 683: 75-84.
21. Druckrey H, Dischler W. (1963) Dosis-Wirkungsbeziehungen bei der Krebszeugung durch 4-dimethylaminostilben bei Ratten. *Z Krebsforsch* 65: 272
22. Druckrey H, Schildbach A, Schmaehl D, Preussmann R, Ivankovic S (1963) Quantitative analysis of the carcinogenic effect of diethylnitrosamine. *Arzneimittelforschung* 13: 841-851.
23. Druckrey H, Schagen B, Ivankovic S (1970) Induction of neurogenic malignancies by one single dose of ethyl-nitrosourea (ENU) given to newborn and juvenile BD IX-strain rats. *Z Krebsforsch* 74: 141-161.
24. Tennekes H, Grettton VA, Stedeford T (2010) Hazard and risk assessment of chemical carcinogenicity within a regulatory context. In: *Cancer risk assessment: Chemical carcinogenesis, hazard evaluation, and risk quantification* (Eds. Hsu CH, Stedeford T), pp. 37-65. John Wiley and Sons, Hoboken, NJ
25. Choe KS, Radparvar F, Matushansky I, Rekhtman N, Han X, Skoultsi AI (2003) Reversal of tumorigenicity and the block to differentiation in erythroleukemia cells by GATA-1. *Cancer Res* 63: 6363-6369.
26. Graf T, Ade N, Beug H (1978) Temperature-sensitive mutant of avian erythroblastosis virus suggests a block of differentiation as mechanism of leukaemogenesis. *Nature* 275: 496-501.
27. Yuspa SH, Kilkenny AE, Stanley J, Lichti U (1985) Keratinocytes blocked in phorbol ester-responsive early stage of terminal differentiation by sarcoma viruses. *Nature* 314: 459-462.
28. Carlborg FW (1981) Dose-response functions in carcinogenesis and the Weibull model. *Food Cosmet Toxicol* 19: 255-263.
29. Sánchez-Bayo F (2009) From simple toxicological models to prediction of toxic effects in time. *Ecotoxicology* 18: 343-354.
30. Beketov MA, Liess M (2008) Acute and delayed effects of the neonicotinoid insecticide thiacloprid on seven freshwater arthropods. *Environ Toxicol Chem* 27: 461-470.
31. Suchail S, Guez D, Belzunces L (2001) Discrepancy between acute and chronic toxicity induced by imidacloprid and its metabolites in *Apis mellifera*. *Environ Toxicol Chem* 20: 2482-2486.
32. Guez D, Suchail S, Gauthier M, Maleszka R, Belzunces LP (2001) Contrasting effects of imidacloprid on habituation in 7- and 8-day-old honeybees (*Apis mellifera*). *Neurobiol Learn Mem* 76: 183-191.
33. Lambin M, Armengaud C, Raymond S, Gauthier M (2001) Imidacloprid-induced facilitation of the proboscis extension reflex habituation in the honeybee. *Arch Insect Biochem Physiol* 48: 129-134.
34. Decourtye A, Devillers J, Cluzeau S, Charreton M, Pham-Delègue M-H (2004) Effects of imidacloprid and deltamethrin on associative learning in honeybees under semi-field and laboratory conditions. *Ecotoxicol Environ Saf* 57: 410-419.
35. Colin ME, Bonmatin JM, Moineau I, Gaimon C, Brun S, et al. (2004) A method to quantify and analyze the foraging activity of honey bees: relevance to sublethal effects induced by systemic insecticides. *Arch Environ Contam Toxicol* 47: 387-395.
36. Chauzat MP, Faucon J-P, Martel A-C, Lachaize J, Cougoule N, et al. (2006) A survey of pesticide residues in pollen loads collected by honey bees in France. *J Econ Entomol* 99:253-262.
37. Yang EC, Chuang YC, Chen YL, Chang LH (2008) Abnormal foraging behavior induced by sublethal dosage of imidacloprid in the honey bee (Hymenoptera: Apidae). *J Econ Entomol* 101: 1743-1748.
38. Decourtye A, Devillers J (2010) Ecotoxicity of neonicotinoid insecticides to bees. In: Thany SH (Ed.) *Insect nicotinic acetylcholine receptors. Advances in Experimental Medicine and Biology* 683: 85-95.
39. Menzel R, Geiger K, Joerges J, Muller ULL, Chittka L (1998) Bees travel novel homeward routes by integrating separately acquired vector memories. *Anim Behav* 55: 139-152.
40. Capaldi EA, Dyer FC (1999) The role of orientation flights on homing performance in honeybees. *J Exp Biol* 202: 1655-1666.
41. Gauthier M (2010) State of the art insect nicotinic acetylcholine receptor function in learning and memory. In: Thany SH (Ed.) *Insect nicotinic acetylcholine receptors. Advances in Experimental Medicine and Biology* 683:97-115.
42. Sánchez-Bayo F, Goka K (2006) Ecological effects of the insecticide imidacloprid and a pollutant from antidandruff shampoo in experimental rice fields. *Environ Toxicol Chem* 25: 1677-1687.
43. Hayasaka D, Korenaga T, Sánchez-Bayo F, Goka K (2011) Differences in ecological impacts of systemic insecticides with different physicochemical properties on biocenosis of experimental paddy fields. *Ecotoxicology DOI* 10.1007/s10646-011-0778-y.
44. Alexander AC, Heard KS, Culp JM (2008) Emergent body size of mayfly survivors *Freshwat Biol* 53: 171-180.
45. Beketov MA, Schäfer RB, Marwitz A, Paschke A, Liess M (2008) Long-term stream invertebrate community alterations induced by the insecticide thiacloprid: Effect concentrations and recovery dynamics. *Sci Total Environ* 405: 96-108.
46. Kreuzweiser DP, Thompson DG, Scarr TA. (2009) Imidacloprid in leaves from systemically treated trees may inhibit litter breakdown by non-target invertebrates. *Ecotoxicol Environ Saf* 72: 1053-1057.
47. Peck DC (2009) Comparative impacts of white grub (Coleoptera: Scarabaeidae) control products on the abundance of non-target soil-active arthropods in turfgrass. *Pedobiologia* 52: 287-299.
48. Sánchez-Bayo F, Yamashita H, Osaka R, Yoneda M, Goka K (2007) Ecological effects of imidacloprid on arthropod communities in and around a vegetable crop. *J Environ Sci Health B* 42: 279-286.
49. Kreuzweiser DP, Good KP, Chartrand DT, Scarr TA, Holmes SB, et al. (2008) Effects on litter-dwelling earthworms and microbial decomposition of soil-applied imidacloprid for control of wood-boring insects. *Pest Manage Sci* 64:112-118.
50. Kreuzweiser DP, Good KP, Chartrand DT, Scarr TA, Thompson DG (2008) Are leaves that fall from imidacloprid-treated maple trees to control Asian longhorned beetles toxic to non-target decomposer organisms? *J Environ Qual* 37: 639-646.
51. Drobne D, Blazic M, Van Gestel CAM, Leser V, Zidar P, et al. (2008) Toxicity of imidacloprid to the terrestrial isopod *Porcellio scaber* (Isopoda, Crustacea). *Chemosphere* 71: 1326 – 1334.
52. Connolly JP (1985) Predicting single-species toxicity in natural water systems. *Environ Toxicol Chem* 4: 573-582.
53. Pascoe D, Shazili NAM (1986) Episodic pollution - A comparison of brief and continuous exposure of rainbow trout to cadmium. *Ecotoxicol Environ Saf* 12: 189-198.
54. Newman MC, Diamond SA, Mulvey M, Dixon P (1989) Allozyme genotype and time to death of mosquitofish, *Gambusia affinis* (Baird and Girard) during acute toxicant exposure: a comparison of arsenate and inorganic mercury. *Aquat Toxicol* 15: 141-156.
55. Mann RM, Vijver MG, Peijnenburg WJGM (2011) Metals and metalloids in terrestrial systems: Bioaccumulation, biomagnification and subsequent adverse effects. In: Sánchez-Bayo, F, van den Brink, PJ, Mann, R (eds.) *Ecological impacts of toxic chemicals*. Bentham Science Publishers pp 43-62.
56. Halstead BW (1959) Minamata disease - a neurological disorder resulting from the eating of Japanese shellfish and other marine organisms. *Amer Malacol Union Ann Rept* 1959: 32
57. Fenlon JS (2002) Time to event analysis in the agricultural sciences. In: Crane, M, Newman, MC, Chapman, PF, Fenlon, J (eds.) *Risk assessment with Time to Events models*. CRC Press LLC, Boca Raton, FL, pp. 103-119.
58. Karman CC (2002) Using time to event modeling to assess the ecological risk of produced water discharges. In: Crane, M, Newman, MC, Chapman, PF,



- Fenlon, J (eds.) Risk assessment with Time to Events models. CRC Press LLC, Boca Raton, FL, pp. 89-102.
59. Kimber A (2002) Time to event analysis in engineering. In: Crane, M, Newman, MC, Chapman, PF, Fenlon, J (eds.) Risk assessment with Time to Events models. CRC Press LLC, Boca Raton, FL, pp. 141-151.
60. Manly BFJ (2002) Time to event analysis in ecology. In: Crane, M, Newman, MC, Chapman, PF, Fenlon, J (eds.) Risk assessment with Time to Events models. CRC Press LLC, Boca Raton, FL, pp. 121-140.
61. Raimondo S, Montague BJ, Barron MG (2007) Determinants of variability in acute to chronic toxicity ratios for aquatic invertebrates and fish. Environ Toxicol Chem 26: 2019-2023.
62. Kenaga EE (1982) Predictability of chronic toxicity from acute toxicity of chemicals in fish and aquatic invertebrates. Environ Toxicol Chem 1: 347-358.
63. Crane M, Chapman PF, Newman MC (2002) Conclusions. In: Crane, M, Newman, MC, Chapman, PF, Fenlon, J (eds.) Risk assessment with Time to Events models. CRC Press LLC, Boca Raton, FL, pp. 167-168.
64. Hoang TC, Tomasso JR, Klaine SJ (2007) An integrated model describing the toxic responses of *Daphnia magna* to pulsed exposures of three metals. Environ Toxicol Chem 26: 132-138.
65. Kooijman SALM (1981) Parametric analyses of mortality rates in bioassays. Water Res 15: 107-119.

This article was originally published in a special issue, **Toxicology of Pesticides** handled by Editor(s). Dr. Francisco Sanchez Bayo, Australia; Dr. Richard ORTEGA, France

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:



- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 200 Open Access Journals
- 15,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: [www.omicsonline.org/submission/](http://www.omicsonline.org/submission/)





**OMICS Publishing Group:**




**Journal of Environmental & Analytical Toxicology:**

Journal of Environmental & Analytical Toxicology is using online manuscript submission, review and tracking systems for quality and quick review processing. Review processing is performed by the editorial board members of Journal of Environmental & Analytical Toxicology and medical informatics or outside experts: at least two independent reviewer's approval followed by editor approval is required for acceptance of any citable manuscript.

The Journal of Environmental & Analytical Toxicology the complicated aspects of addiction, psychological and most probably physical addiction despite of its detrimental consequences and increased tolerance, as deemed by the drug addict himself to his individual health, mental state and unmanageable life style. The Journal is promptly available, and is freely accessible globally through internet to share the innovations of the researchers for scholarly advancement in this field.

Posted 4th April by [OMICS GROUP JOURNALS](#)

0 Add a comment



**OMICS GROUP:**

**Journal of Environmental & Analytical Toxicology:**

Journal of Environmental & Analytical Toxicology is a multidisciplinary field of science concerned with the study of the harmful effects of various chemical, biological and physical agents on individual living organisms.

Journal of Environmental & Analytical Toxicology is an Open Access, peer-reviewed journal which aims to provide the most rapid and reliable source of information on current developments in the field of Environmental Toxicology and Analytical Toxicology. The emphasis will be on publishing quality papers quickly and available to researchers worldwide.

Send feedback

Posted 4th April by [OMICS GROUP JOURNALS](#)



Add a comment



ISSN:2161-0525

# Journal of Environmental & Analytical Toxicology

Open Access

[About Us](#)[Open Access](#)[Conferences](#)[Scientific Credits](#)[Journals](#)[Membership](#)[Submit Manuscript](#)[Register Reviewer / E-Alerts](#)

Search :

Go

[Advanced Search](#)[Home](#)[Join](#)[Contact](#)

## JEAT

- » [Home](#)
- » [Aims and Scope](#)
- » [Editorial Board](#)
- » [Editor-in-Chief](#)
- » [Instructions for Authors](#)
- » [Submit Manuscript](#)
- » [Current Issue](#)
- » [Previous Issue](#)
- » [Special Issues](#)
- » [Articles in Press](#)
- » [Archive](#)
- » [Contact Editorial Office](#)
- » [Indexing and Archiving](#)

[More](#)

### Journal Flier

**Journal of Environmental & Analytical Toxicology**

Open Access



www.omicsonline.org

[Download Journal Flier](#)

### Highly Accessed Articles-2012[Q1]

#### Research Highlights



#### Intertidal Macrofauna Assessment in a Subtropical Mudflat

Two types of sediment cores with different diameters were used to collect sediment samples from an intertidal mudflat to investigate the influence of sampling unit on the quantitative assessment of benthic macroinfaunal communities.

[Prev](#)
[1](#)
[2](#)
[3](#)
[4](#)
[5](#)
[Next](#)

### Environmental & Analytical

#### [Court Greenhouse Gas Decision Potentially](#)

Albany Times Union - Jun 29, 2012  
09-1322, 6/26/12) supporting the Environmental Protection Agency's goal to limit greenhouse gas emissions is not a surprise and is a mixed bag for the U.S. energy economy. Richard Rudden, Chief

[« Previous](#) [Next »](#)[Google Web Elements](#)

### Special Issues

- [Upcoming Special Issues](#)
- [Previous Special Issues](#)

[Submit Manuscript](#)

### Journal of Environmental & Analytical Toxicology

Environmental Toxicology is the study of effects of natural and synthetic pollutants in the environment. Analytical Toxicology deals with the determination of the levels of exposure to potential toxicants via air, water, or food. Journal of Environmental & Analytical Toxicology is concerned with the harmful effects of various chemical, biological, and physical agents on individual living organisms.




### Open Access

Journal of Environmental & Analytical Toxicology is an Open Access, peer-reviewed journal which aims to provide the most rapid and reliable source of information on current developments in the field of Environmental Toxicology and Analytical Toxicology. The emphasis will be on publishing quality papers quickly and freely available to researchers worldwide.



### Biomarkers-2012

3<sup>rd</sup> World Congress on  
**Biomarkers &  
Clinical Research**



July 2-4, 2012  
Embassy Suites Las Vegas,  
USA

### News on

#### Open Access Journals

#### [Free access to British](#)

The Guardian  
The move reflects a groundswell of support for "open access" publishing among academics who have long protested that journal publishers make large profits by locking research behind online